arXiv:0803.0836v3 [hep-th] 21 Apr 2009

KIAS-P08019

Meta-Stable Brane Configurations, Multiple NS5-Branes, and Rotated D6-Branes

Changhyun Ahn

Department of Physics, Kyungpook National University, Taegu 702-701, Korea

`ahn@knu.ac.kr`

Abstract

We construct the type IIA nonsupersymmetric meta-stable brane configurations corresponding to the various $\mathcal{N} = 1$ supersymmetric gauge theories. The D6-branes are both displaced and rotated where these deformations are described as the mass term and quartic term for the fundamental flavors respectively. The multiplicity of the NS5-branes occurs in the superpotential order for adjoint, symmetric, or bifundamental matters. A rich pattern of nonsupersymmetric meta-stable states as well as the supersymmetric stable states is found.

1 Introduction

The dynamical supersymmetry breaking in meta-stable vacua [1, 2] occurs in the $\mathcal{N}=1$ supersymmetric gauge theory with massive fundamental flavors and the corresponding type IIA meta-stable brane constructions have been studied in [3, 4, 5]. Other type IIA nonsupersymmetric meta-stable brane configuration has been constructed by considering the additional quartic term for the quarks in the electric superpotential besides the mass term [6, 7]. This extra deformation in the $\mathcal{N}=1$ supersymmetric gauge theory corresponds to the rotation of D6-branes along the (45)-(89) directions in type IIA string theory. The nonsupersymmetric ground states arise only after the gravitational attraction of NS5-brane [8] is considered.

Branes can be used to describe the dynamics of a wide variety of $\mathcal{N}=1$ supersymmetric gauge theories with different matter contents and superpotential [9]. It is very important to control the gauge singlet meson fields in appropriate way in order to find out the new meta-stable states. In general, one can consider the electric brane configuration of k coincident left NS5-branes(012345) connected by N_c D4-branes to k' coincident right NS5'-branes(012389), with N_f D6-branes(0123789) located between the left NS5- and right NS5'-branes [10]. The creation of D4-branes during the Seiberg dual is related to the gauge singlets in the magnetic brane configuration. Then there exist k magnetic meson fields because each NS5-brane in the above brane configuration produces each meson field when it is crossing the D6-branes.

One can try to find out the new meta-stable brane configuration for a single meson field by taking a single left NS5-brane but multiple k' right NS5'-branes rather than the general case with k left NS5-branes and k' right NS5'-branes where the corresponding dual gauge theory is not known so far. This question is also raised in [11] where the meta-stable brane configuration when $k=2$ and $k'=1$ was studied. In this paper, we find the various meta-stable brane configurations for the different gauge theories with matters and superpotential, along the lines of [12, 13, 14, 15]. All of these examples contain a set of a single NS5-brane and multiple NS5'-branes in their type IIA brane configurations. The single magnetic meson field is realized by the product of quark and anti-quark in the electric gauge theory in the presence of D6-branes with quartic superpotential for the fundamentals. On the other hand, when there are no D6-branes, it is realized by the product of bifundamentals with higher order superpotential.

In section 2, we review the type IIA brane configuration corresponding to the $\mathcal{N}=1$ $SU(N_c)$ gauge theory with two adjoints and fundamentals and deform this theory by adding both the mass term and the quartic term for the fundamentals. Then we describe the dual $\mathcal{N}=1$ $SU(\widetilde{N}_c)$ gauge theory with corresponding dual matter as well as a gauge singlet. We discuss the nonsupersymmetric meta-stable minimum and present the corresponding intersecting brane configuration of type IIA string theory. In section 3, we apply the method of section 2 to the $\mathcal{N}=1$ $Sp(N_c)$ gauge theory with two adjoints and fundamentals.

In section 4, we review the type IIA brane configuration corresponding to the $\mathcal{N}=1$ $SU(N_c)$ gauge theory with symmetric tensor and fundamentals and deform this theory by adding both the mass term and the quartic term for the fundamentals. Then we study the dual $\mathcal{N}=1$ $SU(\widetilde{N}_c)$ gauge theory with corresponding dual matters. We present the nonsupersymmetric meta-stable minimum and the corresponding intersecting brane configuration of type IIA string theory.

In section 5, we review the type IIA brane configuration corresponding to the $\mathcal{N}=1$ $SU(N_c)\times SU(N_c')$ gauge theory with bifundamentals and fundamentals and deform this theory by adding both the mass terms and the quartic terms for each fundamentals. Then we describe the dual $\mathcal{N}=1$ $SU(\widetilde{N}_c)\times SU(N_c')$ gauge theory with corresponding dual matters. We discuss the nonsupersymmetric meta-stable minimum and the corresponding intersecting brane configuration. In section 6, we use the method of section 5 to the $\mathcal{N}=1$ $Sp(N_c)\times SO(2N_c')$ gauge theory with bifundamentals, vectors and fundamentals. In section 7, we apply the method of section 5 to the $\mathcal{N}=1$ $SU(N_c)\times SO(2N_c')$ gauge theory with bifundamentals, vectors and fundamentals.

In section 8, we review the type IIA brane configuration corresponding to the $\mathcal{N}=1$ $SU(N_c)\times SU(N_c')$ gauge theory with symmetric tensor and bifundamentals and deform this theory by adding both the mass term and the higher order term for bifundamentals. Then we describe the dual $\mathcal{N}=1$ $SU(\widetilde{N}_c)\times SU(N_c')$ gauge theory with corresponding dual matters. We discuss the nonsupersymmetric meta-stable minimum and the corresponding intersecting brane configuration. In section 9, we use the method of section 8 to the $\mathcal{N}=1$ $SU(N_c)\times SU(N_c')$ gauge theory with antisymmetric tensor, eight fundamentals and bifundamentals.

In section 10, we review the type IIA brane configuration corresponding to the $\mathcal{N}=1$ $SU(N_c)\times SU(N_c')$ gauge theory with symmetric tensor, bifundamentals and fundamentals and deform this theory by adding both the mass terms and the quartic terms for each fundamentals. Then we describe the dual $\mathcal{N}=1$ $SU(\widetilde{N}_c)\times SU(N_c')$ gauge theory or $\mathcal{N}=1$ $SU(N_c)\times SU(\widetilde{N}_c')$ gauge theory with corresponding dual matters. We study the nonsupersymmetric meta-stable minimum and the corresponding intersecting brane configurations. In section 11, we also describe the $\mathcal{N}=1$ $SU(N_c)\times SU(N_c')$ gauge theory with antisymmetric tensor, bifundamentals and fundamentals.

In section 12, we make some comments for the future directions.

2 $SU(N_c)$ with two adj. and N_f-fund.

2.1 Electric theory

The type IIA supersymmetric electric brane configuration [10] corresponding to $\mathcal{N} = 1$ $SU(N_c)$ gauge theory with two adjoint fields Φ, Φ' and N_f-fundamental flavors $Q, \widetilde{Q}$ can be described as follows: one NS5-brane(012345), k' NS5'-branes(012389), N_c D4-branes(01236), and N_f D6-branes(0123789). Let us introduce two complex coordinates [9]

$$v \equiv x^4 + ix^5, \qquad w \equiv x^8 + ix^9.$$

Since we consider a single NS5-brane, the adjoint field Φ is massive and can be integrated out. For large coupling in front of the quadratic Φ term, there is no Φ-dependence in the superpotential. The N_c-color D4-branes are suspended between the left NS5-brane and the right NS5'-branes and similarly the N_f D6-branes are located between the left NS5-brane and the right NS5'-branes. The N_f-fundamental flavors $Q, \widetilde{Q}$ are strings stretching between N_f D6-branes and N_c-color D4-branes while the adjoint field Φ' is related to the fluctuations of N_c-color D4-branes in v direction.

Let us deform this theory by adding the mass term and the quartic term for fundamental quarks $Q, \widetilde{Q}$ in order to find the new nonsupersymmetric meta-stable states. The former can be achieved by displacing the D6-branes along $+v$ direction leading to their coordinates $v = +v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote those rotated D6-branes by $D6_{-\theta}$-branes. Then, in the electric gauge theory, the deformed superpotential is given by

$$W_{elec} = \left[\frac{g_{\Phi'}}{2} \operatorname{tr} \Phi'^{k'+1} + Q\Phi'\widetilde{Q}\right] - m \operatorname{tr} Q\widetilde{Q} + \frac{\alpha}{2} \operatorname{tr}(Q\widetilde{Q})^2, \qquad \alpha = \frac{\tan\theta}{\Lambda}, \qquad m = \frac{v_{D6_{-\theta}}}{2\pi\ell_s^2} \tag{2.1}$$

where Λ is related to the scales of the electric and magnetic theories and $+v_{D6_{-\theta}}$ is the v coordinate of the center of coincident rotated $D6_{-\theta}$-branes. When $k' = 1$, this theory reduces to the one in [6, 7] because the first two terms of (2.1) contribute to the additional quartic term for the quarks $Q, \widetilde{Q}$ and we are left with the last two terms in (2.1). Therefore, we focus on the nontrivial case with the number of NS5'-branes $k' \geq 2$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration [1] with superpotential (2.1) in type IIA string theory as follows and draw this in Figure 1:

---

[1] Without the quartic term($\alpha = 0$), the theory with $k' = N_c - 1$ obtained by resolving the superpotential(i.e., there exist also the lower order terms by splitting the k' NS5'-branes in v direction) was studied in [16] some time ago in the supersymmetric brane configuration and this theory without D6-branes when $k' \geq N_c + 1$(multitrace interactions) was also studied in the context of supersymmetry breaking vacua from M5-branes recently [17].

- One left NS5-brane in (012345) directions with $w = 0$
- k' right NS5'-branes in (012389) directions with $v = 0$
- N_c-color D4-branes in (01236) directions with $v = 0 = w$
- N_f $D6_{-\theta}$-branes in (01237) directions and two other directions in (v, w)-plane



Figure 1: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c)$ with two adjoints and fundamentals $Q, \widetilde{Q}$. Note that there are k' NS5'-branes and the fluctuations of N_c color D4-branes in v direction correspond to the adjoint field. A "rotation" of N_f D6-branes in (w, v)-plane, which become $D6_{-\theta}$-branes, corresponds to a quartic term for the fundamentals while a "displacement" of N_f D6-branes in $+v$ direction corresponds to a mass term for the fundamentals.

2.2 Magnetic theory

Let us move the left NS5-brane to the right all the way past the right NS5'-branes and one arrives at the Figure 2A. Note that there exists a creation of N_f-flavor D4-branes connecting N_f $D6_{-\theta}$-branes and k' NS5'-branes. Recall that the N_f $D6_{-\theta}$-branes are not parallel to the NS5-brane in Figure 1 unless $\theta = \frac{\pi}{2}$. The linking number [18] of the NS5-brane from Figure 2A is $l_m = \frac{N_f}{2} - \widetilde{N}_c$. On the other hand, the linking number of the NS5-brane from Figure 1 is $l_e = -\frac{N_f}{2} + N_c$. From the equality between these two relations, one obtains the number of colors of dual magnetic theory [10]

$$\widetilde{N}_c = N_f - N_c.$$

Note that the multiplicity k' of NS5'-branes does not arise in this computation because the creation of flavor D4-branes appears via the dual process between the NS5-brane and N_f $D6_{-\theta}$-branes which do not depend on k'.

The low energy theory on the dual color D4-branes has $SU(\widetilde{N}_c)$ gauge group and an adjoint field ϕ' coming from 4-4 strings connecting the dual color D4-branes and N_f-fundamental dual quarks $q, \widetilde{q}$ coming from 4-4 strings connecting between the color D4-branes and flavor D4-branes. Moreover, a single magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from



Figure 2: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 1 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(2A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(2B). The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into $\widetilde{N}_c$-, $(N_c - l)$- and l- D4-branes(2A). Further n- D4-branes among $(N_c - l)$- D4-branes are moved to the NS5-brane(2B).

4-4 strings of flavor D4-branes. Then the dual magnetic superpotential is given by

$$W_{mag} = \left[\frac{g_{\phi'}}{2} \operatorname{tr} {\phi'}^{k'+1} + q \phi' \widetilde{q} + \frac{1}{\Lambda} M q \widetilde{q} \right] + \frac{\alpha}{2} \operatorname{tr} M^2 - m M. \tag{2.2}$$

The undeformed expression, the first three terms, is already found in [10]. The case with $k' = 1$ leads to the one in [6]. In order to obtain the supersymmetric vacua, one computes the F-term equations for the superpotential (2.2):

$$\begin{aligned} \frac{1}{\Lambda} M q + q \phi' &= 0, \qquad \phi' \widetilde{q} + \frac{1}{\Lambda} \widetilde{q} M = 0, \\ \frac{1}{2} g_{\phi'} (k'+1) {\phi'}^{k'} + \widetilde{q} q &= 0, \qquad \frac{1}{\Lambda} q \widetilde{q} = m - \alpha M. \end{aligned} \tag{2.3}$$

We choose the adjoint field ϕ' to be diagonal, i.e., $\phi' = \operatorname{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c})$. Then the first two equations of (2.3) imply that the upper left $\widetilde{N}_c \times \widetilde{N}_c$ block of $\frac{M}{\Lambda}$ is given by $\operatorname{diag}(-\phi'_1, \cdots, -\phi'_{\widetilde{N}_c})$. By substituting this into the last equation of (2.3), one obtains the upper left $\widetilde{N}_c \times \widetilde{N}_c$ block of $\frac{1}{\Lambda} q \widetilde{q}$ is given by $\operatorname{diag}(m + \alpha \Lambda \phi'_1, \cdots, m + \alpha \Lambda \phi'_{\widetilde{N}_c})$. Moreover, the lower right $(N_f - \widetilde{N}_c) \times (N_f - \widetilde{N}_c)$ block of M is given by l's zero eigenvalues and $(N_f - \widetilde{N}_c - l)$'s eigenvalues $\frac{m}{\alpha}$. Then finally, the third equation of (2.3) gives rise to the expectation value for the adjoint field ϕ' satisfying that

$$\frac{1}{2} g_{\phi'} (k'+1) {\phi'_j}^{k'} = -\Lambda (m + \alpha \Lambda \phi'_j) \tag{2.4}$$

for $j = 1, 2, \cdots, \widetilde{N}_c$. For nonzero quark mass m, the expectation value ϕ'_j is not vanishing and $-\Lambda \phi'_j > \frac{m}{\alpha}$ for the positiveness of left hand side of (2.4) when $g_{\phi'}$ and ϕ'_j are real.

Let us first describe the nonsupersymmetric meta-stable states and supersymmetric ones when all the N_f $D6_{-\theta}$-branes and k' NS5'-branes are coincident with each other.

- Coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes(equal massive flavors)

One writes $N_f \times N_f$ matrix M with $\widetilde{N}_c(= N_f - N_c)$ eigenvalues by the diagonal elements for ϕ', l's eigenvalues by the zeros and $(N_c - l)$ eigenvalues $\frac{m}{\alpha}$ as follows:

$$M = \left(\begin{array}{ccc} -\Lambda\phi' & 0 & 0 \\ 0 & 0_l & 0 \\ 0 & 0 & \frac{m}{\alpha}\mathbf{1}_{N_c-l} \end{array} \right), \qquad \text{with} \qquad \phi' = \text{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c}). \tag{2.5}$$

In the brane configuration of Figure 2A, the l of the N_f-flavor D4-branes are connected with l-color D4-branes and the resulting l D4-branes stretch from the $D6_{-\theta}$-branes to the NS5-brane directly and the intersection point between the l D4-branes and the $D6_{-\theta}$-branes is given by $(v, w) = (+v_{D6_{-\theta}}, 0)$. This corresponds to exactly the l's eigenvalues from zeros of M in (2.5). Now the remaining $(N_c - l)$-flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes correspond to the eigenvalues of M in (2.5), i.e., $\frac{m}{\alpha}\mathbf{1}_{N_c-l}$. The intersection point between the $(N_c - l)$ D4-branes and the NS5'-branes is given by $(v, w) = (0, +v_{D6_{-\theta}} \cot\theta)$ from trigonometric geometry [6]. Finally, the remnant $\widetilde{N}_c$-flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes correspond to the eigenvalues $-\Lambda\phi'$ in (2.5) with (2.4) providing the exact nonzero w coordinates for these flavor D4-branes.

One also represents the vacuum expectation value for the quarks as follows:

$$q\widetilde{q} = \left(\begin{array}{ccc} \Lambda(m\mathbf{1}_{\widetilde{N}_c} + \alpha\Lambda\phi') & 0 & 0 \\ 0 & \Lambda m\mathbf{1}_l & 0 \\ 0 & 0 & 0_{N_c-l} \end{array} \right), \qquad \text{with} \qquad \phi' = \text{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c}). \tag{2.6}$$

In the l-th vacuum the gauge symmetry is broken to $SU(\widetilde{N}_c - l)$ and when the supersymmetric vacuum is drawn in Figure 2A with $l = 0$, the gauge group $SU(\widetilde{N}_c)$ is unbroken. Then the supersymmetric ground state corresponds to the vacuum expectation values M by $-\Lambda\phi'$ which has $\widetilde{N}_c$'s eigenvalues and $\frac{m}{\alpha}$ with degeneracy N_c.

The theory has nonsupersymmetric meta-stable ground states since there exists an attractive gravitational interaction between the flavor D4-branes and the NS5-brane from the DBI action [8]. Let us rescale the meson field as $M = h\Lambda\Phi$. Then the magnetic superpotential (2.2) can be rewritten in terms of $\Phi, q, \widetilde{q}$ and ϕ'

$$W_{dual} = h\Phi q\widetilde{q} + \frac{\mu_\phi}{2} h^2 \,\text{tr}\, \Phi^2 - h\mu^2 \,\text{tr}\, \Phi + \frac{g_{\phi'}}{2} \,\text{tr}\, \phi'^{k'+1} + q\phi'\widetilde{q} \tag{2.7}$$

with $\mu^2 = m\Lambda$ and $\mu_\phi = \alpha\Lambda^2$.

Now one splits the $(N_c - l) \times (N_c - l)$ block at the lower right corner of M and $q\widetilde{q}$ into blocks of size n and $(N_c - l - n)$ and then (2.5) and (2.6) are rewritten as follows [7]:

$$
\begin{aligned}
h\Phi &= \begin{pmatrix} -\phi' & 0 & 0 & 0 \\ 0 & 0_l & 0 & 0 \\ 0 & 0 & h\Phi_n & 0 \\ 0 & 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{N_c-l-n} \end{pmatrix}, \\
q\widetilde{q} &= \begin{pmatrix} \mu^2\mathbf{1}_{\widetilde{N}_c} + \alpha\Lambda^2\phi' & 0 & 0 & 0 \\ 0 & \mu^2\mathbf{1}_l & 0 & 0 \\ 0 & 0 & \varphi\widetilde{\varphi} & 0 \\ 0 & 0 & 0 & 0_{N_c-l-n} \end{pmatrix}
\end{aligned}
\tag{2.8}
$$

with $\phi' = \mathrm{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c})$. Here φ and $\widetilde{\varphi}$ are $n \times (\widetilde{N}_c - l)$ matrices and correspond to n-flavors of fundamentals of the gauge group $SU(\widetilde{N}_c - l)$ which is unbroken. One can move n D4-branes from $(N_c - l)$ flavor D4-branes stretched between the $D6_{-\theta}$-branes and the NS5'-branes at $w = +v_{D6_{-\theta}} \cot\theta$, to the local minimum of the potential and the end points of these n D4-branes are at a nonzero w [6]. In the brane configuration from Figure 2B, φ and $\widetilde{\varphi}$ correspond to fundamental strings connecting between the n-flavor D4-branes and $(\widetilde{N}_c - l)$-color D4-branes. Moreover, the $h\Phi_n$ and $\varphi\widetilde{\varphi}$ are $n \times n$ matrices. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_n = \frac{\mu^2}{\mu_\phi}\mathbf{1}_n$ and $\varphi\widetilde{\varphi} = 0$.

The full one loop potential from (2.7) and (2.8) takes the form

$$
\frac{V}{|h|^2} = |\Phi_n\varphi + \varphi\phi'|^2 + |\widetilde{\varphi}\Phi_n + \phi'\widetilde{\varphi}|^2 + \left|\varphi\widetilde{\varphi} - \mu^2\mathbf{1}_n + h\mu_\phi\Phi_n\right|^2 + b|h\mu|^2 \,\mathrm{tr}\, \Phi_n^\dagger \Phi_n \tag{2.9}
$$

where $b = \frac{(\ln 4 - 1)}{8\pi^2}\widetilde{N}_c$ and we did not include Φ_n- or $\Phi_n^\dagger$-independent term and after differentiating this (2.9) with respect to $\Phi_n^\dagger$ one obtains the local nonzero stable point given by

$$
h\Phi_n \simeq \frac{\mu_\phi}{b}\mathbf{1}_n \qquad \text{or} \qquad M_n \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_c}\mathbf{1}_n .
$$

This corresponds to the location of w coordinate, which is less than $\frac{m}{\alpha\Lambda}(= \frac{\mu^2}{\mu_\phi})$, of n flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes.

- Non-coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes(different massive flavors)

Let us suppose that the numbers of $D6_{-\theta}$-branes and the NS5'-branes are equal to each other:$N_f = k'$. Let us displace the k' $D6_{-\theta}$-branes and NS5'-branes given in Figure 2A in the v direction respectively to two k' different points denoted by $v_{D6_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$. The color and flavor D4-branes attached to them are displaced also. The number of color D4-branes stretched between j-th $NS5'_j$-brane and the NS5-brane is denoted by $\widetilde{N}_{c,j}$ while the number of flavor D4-branes stretched between the j-th $D6_{-\theta,j}$-brane and the j-th $NS5'_j$-brane is denoted by $N_{f,j}$. Then it is obvious that there are relations between the color and flavor D4-branes $\sum_{j=1}^{k'} \widetilde{N}_{c,j} = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N_{f,j} = N_f$. When all the $NS5'_j$-branes and $D6_{-\theta,j}$-branes are distinct, the low energy physics corresponds to k'-decoupled supersymmetric gauge theories with gauge group $\prod_{j=1}^{k'} SU(\widetilde{N}_{c,j})$.

One deforms the Figure 2B by displacing the multiple $D6_{-\theta}$-branes and $NS5'$-branes along v direction, as in [15]. Then the n curved flavor D4-branes attached to them(as well as other D4-branes) are displaced also as k' different n_j's connecting between $D6_{-\theta,j}$-brane and $NS5'_j$-brane. When we rescale the submeson field as $M_j = h\Lambda\Phi_j$ [15], then the Kahler potential for Φ_j is canonical and the magnetic quarks q_j and $\widetilde{q}_j$ are canonical near the origin of field space [1]. Then the magnetic superpotential (2.7) can be rewritten in terms of $\Phi_j, q_j, \widetilde{q}_j$ and $\phi'_{\widetilde{N}_{c,j}}$ which is coming from 4-4 strings connecting between the j-th $\widetilde{N}_{c,j}$ D4-branes

$$W_{dual} = \sum_{j=1}^{k'} \left[h\Phi_j q_j \widetilde{q}_j + \frac{\mu_\phi}{2} h^2 \operatorname{tr} \Phi_j^2 - h \operatorname{tr} \mu_j^2 \Phi_j + \frac{g_{\phi'}}{2} \operatorname{tr} {\phi'}_{\widetilde{N}_{c,j}}^{k'+1} + q_j \phi'_{\widetilde{N}_{c,j}} \widetilde{q}_j \right] \tag{2.10}$$

with $\mu_j^2 = m_j \Lambda_j$ and $\mu_\phi = \alpha\Lambda^2$ as before.

One splits the $(N_{c,j} - l_j) \times (N_{c,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $q_j\widetilde{q}_j$ into blocks of size n_j and $(N_{c,j} - l_j - n_j)$ for all j as follows [15]:

$$\begin{aligned} h\Phi &= \left(\begin{array}{ccccc} 0_{\widetilde{N}_c+l+n} & 0 & 0 & \cdots & 0 \\ 0 & \frac{\mu_1^2}{\mu_\phi}\mathbf{1}_{N_{c,1}-l_1-n_1} & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \frac{\mu_{k'}^2}{\mu_\phi}\mathbf{1}_{N_{c,k'}-l_{k'}-n_{k'}} \end{array} \right) \\ &- \operatorname{diag}(\phi'_{\widetilde{N}_{c,1}}, \cdots, \phi'_{\widetilde{N}_{c,k'}}, 0_{N_c}) + \operatorname{diag}(0_{\widetilde{N}_c+l}, h\Phi_{n_1}, \cdots, h\Phi_{n_{k'}}, 0_{N_c-l-n}) \end{aligned} \tag{2.11}$$

and

$$\begin{aligned} q\widetilde{q} &= \left(\begin{array}{ccccccc} 0_{\widetilde{N}_c} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \mu_1^2\mathbf{1}_{l_1} & 0 & 0 & \cdots & 0 & 0 \\ 0 & \cdot & \cdot & \cdot & \cdots & \cdot & 0 \\ 0 & 0 & 0 & 0 & \cdots & \mu_{k'}^2\mathbf{1}_{l_{k'}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0_{N_c-l} \end{array} \right) \\ &+ \operatorname{diag}(\mu_1^2\mathbf{1}_{\widetilde{N}_{c,1}} + \alpha\Lambda^2\phi'_{\widetilde{N}_{c,1}}, \cdots, \mu_{k'}^2\mathbf{1}_{\widetilde{N}_{c,k'}} + \alpha\Lambda^2\phi'_{\widetilde{N}_{c,k'}}, 0_{N_c}) \\ &+ \operatorname{diag}(0_{\widetilde{N}_c+l}, \varphi_{n_1}\widetilde{\varphi}_{n_1}, \cdots, \varphi_{n_{k'}}\widetilde{\varphi}_{n_{k'}}, 0_{N_c-l-n}) \end{aligned} \tag{2.12}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$. Here φ_{n_j} and $\widetilde{\varphi}_{n_j}$ are $n_j \times (\widetilde{N}_{c,j} - l_j)$ matrices and correspond to n_j-flavors of fundamentals of the gauge group $SU(\widetilde{N}_{c,j} - l_j)$ which is unbroken. They correspond to fundamental strings connecting between the n_j-flavor D4-branes and $(\widetilde{N}_{c,j} - l_j)$-color D4-branes. Moreover, the Φ_{n_j} and $\varphi_{n_j}\widetilde{\varphi}_{n_j}$ are $n_j \times n_j$ matrices. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{n_j} = \frac{\mu_j^2}{\mu_\phi}\mathbf{1}_{n_j}$ and $\varphi_{n_j}\widetilde{\varphi}_{n_j} = 0$. The full one loop potential from (2.10), (2.11) and (2.12) can be written similarly and the local nonzero stable point arises as

$$h\Phi_{n_j} \simeq \frac{\mu_\phi}{b_j}\mathbf{1}_{n_j} \qquad \text{or} \qquad M_{n_j} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_{c,j}}\mathbf{1}_{n_j}$$

corresponding to the nonzero w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane.

Therefore, the meta-stable states, for fixed k' which is related to the order of the adjoint field in the superpotential and θ which is a deformation parameter by rotation angle of $D6_{-\theta}$-branes, are classified by the number of various D4-branes and the positions of multiple $D6_{-\theta}$-branes and NS5'-branes: $(N_{c,j}, N_{f,j}, l_j, n_j)$ and $(v_{D6_{-\theta,j}}, v_{NS5'_j})$ where $j = 1, 2, \cdots, k'$.

When N_f is not equal to k'(for example, when $N_f > k'$), then some coincident $D6_{-\theta}$-branes among N_f $D6_{-\theta}$-branes should be connected to the NS5'-branes in order to connect all the flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes. These coincident $D6_{-\theta}$-branes can be obtained by taking those quark masses equal. Then all the previous descriptions for the meta-stable states for the case with $N_f = k'$ can be applied in this case also without any difficulty. On the other hand, when $N_f < k'$, then some coincident NS5'-branes among k' NS5'-branes should be connected to the $D6_{-\theta}$-branes.

3 $Sp(N_c)$ with two adj. and N_f-fund.

3.1 Electric theory

The type IIA supersymmetric electric brane configuration [10, 19, 20] corresponding to $\mathcal{N} = 1$ $Sp(N_c)$ gauge theory with two adjoint fields Φ, Φ'(which are symmetric tensors in the symplectic gauge group) and N_f-fundamental flavors Q can be described as follows: one NS5-brane, $(2k'+1)$ NS5'-branes, $2N_c$ D4-branes, $2N_f$ D6-branes and the $O4^{\pm}$-planes(01236). The $2N_c$-color D4-branes are suspended between the left NS5-brane and the right NS5'-branes and the $2N_f$ D6-branes are located between the left NS5-brane and the right NS5'-branes. The $2N_f$-fields Q are strings stretching between $2N_f$ D6-branes and $2N_c$-color D4-branes. The adjoint field Φ' is related to the fluctuations of color D4-branes in v direction.

Let us deform this theory by adding the mass and the quartic terms for fundamental quarks. The former can be achieved by displacing the D6-branes along $\pm v$ direction leading to their coordinates $v = \pm v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $D6_{-\theta}$-branes. That is, each N_f

D6-branes is moving to the $\pm v$ directions respectively and then is rotating by an angle $-\theta$ in (w, v)-plane. Therefore, in the electric gauge theory, the deformed superpotential is given by

$$W_{elec} = \left[\frac{g_{\Phi'}}{2}\,\mathrm{tr}\,\Phi'^{2k'+2} + Q\Phi' Q\right] - m\,\mathrm{tr}\,QQ + \frac{\alpha}{2}\,\mathrm{tr}(QQ)^2, \qquad \alpha = \frac{\tan\theta}{\Lambda}, \quad m = \frac{v_{D6_{-\theta}}}{2\pi\ell_s^2}. \quad (3.1)$$

When $k' = 0$, this theory reduces to the one similar to the unitary case [6, 7] because the first two terms of (3.1) contribute to the additional quartic term for the quarks. Therefore, we focus on the nontrivial case with $k' \geq 1$. The order of the first term of (3.1) can be determined by replacing the k' of (2.1) with $(2k'+1)$ in the sense that when the additional k' NS5'-branes are added into a single NS5'-brane, their mirrors also should be present, leading to the total number of NS5'-branes being $(2k'+1)$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (3.1) in type IIA string theory as follows and draw this in Figure 3 which is nothing but the Figure 1 together with the addition of $O4^{\pm}$-planes:

- One left NS5-brane in (012345) directions with $w = 0$
- $(2k'+1)$ right NS5'-branes in (012389) directions with $v = 0$
- $2N_c$-color D4-branes in (01236) directions with $v = 0 = w$
- $2N_f$ $D6_{-\theta}$-branes in (01237) directions and two other directions in (v, w)-plane
- $O4^{\pm}$-planes in (01236) directions with $v = 0 = w$



Figure 3: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $Sp(N_c)$ with two adjoints and fundamentals Q. Note that there are $(2k'+1)$ NS5'-branes and the fluctuations of $2N_c$ color D4-branes in v direction correspond to the adjoint field. A rotation of N_f D6-branes in (w, v)-plane corresponds to a quartic term for the fundamentals while a displacement of N_f D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals.

3.2 Magnetic theory

Let us move the left NS5-brane to the right past the right NS5'-branes as in previous section and we arrive at the Figure 4A. Note that there exists a creation of N_f-flavor D4-branes connecting N_f $D6_{-\theta}$-branes and $(2k'+1)$ NS5'-branes(and their mirrors). The linking number of the NS5-brane from Figure 4A is $l_m = \frac{2N_f}{2} - 2 - 2\widetilde{N}_c$. The linking number of the NS5-brane from Figure 3 is $l_e = -\frac{2N_f}{2} + 2 + 2N_c$. From the equality between these, one obtains the number of colors of dual magnetic theory [10, 1, 21] $\widetilde{N}_c = N_f - N_c - 2$.



Figure 4: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration corresponding to Figure 3 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(4A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(4B). The N_f upper flavor D4-branes connecting between upper $D6_{-\theta}$-branes and NS5'-branes are splitting into $\widetilde{N}_c$-, $(N_f - \widetilde{N}_c - l)$- and l- D4-branes(4A). Further n- D4-branes among upper $(N_f - \widetilde{N}_c - l)$- D4-branes are moved to the NS5-brane(4B).

The low energy theory on the color D4-branes has $Sp(\widetilde{N}_c)$ gauge group and an adjoint field ϕ' coming from 4-4 strings connecting the color D4-branes and N_f-fundamental quarks q coming from 4-4 strings connecting between the color D4-branes and flavor D4-branes. Moreover, a single magnetic meson field $M \equiv QQ$ is $2N_f \times 2N_f$ matrix and comes from 4-4 strings of flavor D4-branes. Then the magnetic superpotential is given by

$$W_{mag} = \left[\frac{g_{\phi'}}{2} \operatorname{tr} \phi'^{2k'+2} + q\phi' q + \frac{1}{\Lambda} M q q \right] + \frac{\alpha}{2} \operatorname{tr} M^2 - mM. \tag{3.2}$$

When $k' = 0$ and there is no mass term for M, this superpotential reduces to the one found in [1, 21].

In order to obtain the supersymmetric vacua, one computes the F-term equations for the superpotential (3.2):

$$\begin{aligned} \frac{1}{\Lambda} M q + q\phi' &= 0, \qquad g_{\phi'}(k'+1)\phi'^{2k'+1} + qq = 0, \\ \frac{1}{\Lambda} qq &= m - \alpha M. \end{aligned} \tag{3.3}$$

We choose the adjoint field ϕ' to be diagonal, i.e., $\phi' = \operatorname{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c}) \otimes \sigma_3$ [19]. Then the first equation of (3.3) implies that the upper left $2\widetilde{N}_c \times 2\widetilde{N}_c$ block of $\frac{M}{\Lambda}$ is given by

$\mathrm{diag}(-\phi'_1, \cdots, -\phi'_{\widetilde{N}_c}) \otimes i\sigma_2$. By substituting this into the last equation of (3.3), one obtains the upper left $2\widetilde{N}_c \times 2\widetilde{N}_c$ block of $\frac{1}{\Lambda} qq$ is given by $\mathrm{diag}(m + \alpha\Lambda\phi'_1, \cdots, m + \alpha\Lambda\phi'_{\widetilde{N}_c}) \otimes i\sigma_2$. Moreover, the lower right $2(N_f - \widetilde{N}_c) \times 2(N_f - \widetilde{N}_c)$ block of M is given by $2l$'s zero eigenvalues and $2(N_f - \widetilde{N}_c - l)$'s eigenvalues $\pm\frac{m}{\alpha}$. Then finally, the second equation of (3.3) gives rise to the expectation value for the adjoint field ϕ' satisfying that

$$g_{\phi'}(k'+1){\phi'_j}^{2k'+1} = -\Lambda(m + \alpha\Lambda\phi'_j) \tag{3.4}$$

for $j = 1, 2, \cdots, \widetilde{N}_c$. For nonzero quark mass m, the expectation value ϕ'_j is not vanishing and $-\Lambda\phi'_j > \frac{m}{\alpha}$ for the positiveness of left hand side of (3.4) when $g_{\phi'}$ and ϕ'_j are real.

Let us first describe the nonsupersymmetric meta-stable states and supersymmetric ones when all the $2N_f$ $D6_{-\theta}$-branes and $(2k'+1)$ NS5'-branes are coincident with each other.

- Two coincident N_f $D6_{-\theta}$-branes and coincident $(2k'+1)$ NS5'-branes

One writes $2N_f \times 2N_f$ matrix M with $2\widetilde{N}_c$ eigenvalues by the diagonal elements for $\pm\varphi'$, $2l$'s eigenvalues by the zeros and $2(N_f - \widetilde{N}_c - l)$ eigenvalues $\pm\frac{m}{\alpha}$ as follows:

$$M = \begin{pmatrix} -\Lambda\varphi' \otimes i\sigma_2 & 0 & 0 \\ 0 & 0_{2l} & 0 \\ 0 & 0 & \frac{m}{\alpha}\mathbf{1}_{N_f - \widetilde{N}_c - l} \otimes i\sigma_2 \end{pmatrix}, \quad \varphi' = \mathrm{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c}). \tag{3.5}$$

Therefore, in the brane configuration of Figure 4A, the l of the upper N_f flavor D4-branes are connected with l of $\widetilde{N}_c$ color D4-branes and the resulting D4-branes stretch from the upper $D6_{-\theta}$-branes to the NS5-brane directly and the intersection point between the l upper D4-branes and the NS5-brane is given by $(v, w) = (+v_{D6_{-\theta}}, 0)$. Now the $(N_f - \widetilde{N}_c - l)$ upper flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes are related to the corresponding half eigenvalues of M:$\frac{m}{\alpha}\mathbf{1}_{N_f - \widetilde{N}_c - l}$. The intersection point between the $(N_f - \widetilde{N}_c - l)$ upper D4-branes and the NS5'-branes is given by $(v, w) = (0, +v_{D6_{-\theta}} \cot\theta)$ corresponding to positive eigenvalues of M. Finally, the remnant $2\widetilde{N}_c$-flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes correspond to the eigenvalues $\pm\varphi'$ in (3.5) with (3.4) which shows the exact coordinate of w for these flavor D4-branes.

One represents the vacuum expectation value for the quarks as follows:

$$qq = \begin{pmatrix} \Lambda(m\mathbf{1}_{\widetilde{N}_c} + \alpha\Lambda\varphi') \otimes i\sigma_2 & 0 & 0 \\ 0 & \Lambda m \mathbf{1}_l \otimes i\sigma_2 & 0 \\ 0 & 0 & 0_{2(N_f - \widetilde{N}_c - l)} \end{pmatrix}, \varphi' = \mathrm{diag}(\phi'_1, \cdots, \phi'_{\widetilde{N}_c}). \tag{3.6}$$

In the l-th vacuum the gauge symmetry is broken to $Sp(\widetilde{N}_c - l)$ and when the supersymmetric vacuum is drawn in Figure 4A with $l = 0$, the gauge group $Sp(\widetilde{N}_c)$ is unbroken. Then the supersymmetric ground state corresponds to the vacuum expectation values M by $\pm\varphi'$ which has $2\widetilde{N}_c$'s eigenvalues and $\pm\frac{m}{\alpha}$ with degeneracy $2(N_f-\widetilde{N}_c)$.

Let us rescale the meson field as $M = h\Lambda\Phi$ [1]. Then the magnetic superpotential (3.2) can be rewritten in terms of Φ, q and ϕ'

$$W_{dual} = h\Phi qq + \frac{\mu_\phi}{2}h^2\,\mathrm{tr}\,\Phi^2 - h\mu^2\,\mathrm{tr}\,\Phi + \frac{g_{\phi'}}{2}\,\mathrm{tr}\,\phi'^{2k'+2} + q\phi' q \tag{3.7}$$

where $\mu^2 = m\Lambda$ and $\mu_\phi = \alpha\Lambda^2$.

Now one splits the $2(N_f-\widetilde{N}_c-l)\times 2(N_f-\widetilde{N}_c-l)$ block at the lower right corner of M and qq into blocks of size $2n$ and $2(N_f-\widetilde{N}_c-l-n)$ and then (3.5) and (3.6) are rewritten as follows:

$$\begin{aligned} h\Phi &= \left(\begin{array}{cccc} -\varphi'\otimes i\sigma_2 & 0 & 0 & 0 \\ 0 & 0_{2l} & 0 & 0 \\ 0 & 0 & h\Phi_{2n} & 0 \\ 0 & 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{(N_f-\widetilde{N}_c-l-n)}\otimes i\sigma_2 \end{array}\right), \\ qq &= \left(\begin{array}{cccc} (\mu^2\mathbf{1}_{\widetilde{N}_c}+\alpha\Lambda^2\varphi')\otimes i\sigma_2 & 0 & 0 & 0 \\ 0 & \mu^2\mathbf{1}_l\otimes i\sigma_2 & 0 & 0 \\ 0 & 0 & \varphi\varphi & 0 \\ 0 & 0 & 0 & 0_{2(N_f-\widetilde{N}_c-l-n)} \end{array}\right). \end{aligned}$$

Here φ is $2n\times 2(\widetilde{N}_c-l)$ dimensional matrix and corresponds to $2n$ flavors of fundamentals of the gauge group $Sp(\widetilde{N}_c-l)$ which is unbroken by the nonzero expectation value of q. The Φ_{2n} and $\varphi\varphi$ are $2n\times 2n$ matrices. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{2n} = \frac{\mu^2}{\mu_\phi}\mathbf{1}_n\otimes i\sigma_2$ and $\varphi = 0$.

The full one loop potential takes the form

$$\frac{V}{|h|^2} = |\Phi_{2n}\varphi+\varphi\varphi'|^2 + \left|\varphi\varphi-\mu^2\mathbf{1}_{2n}+h\mu_\phi\Phi_{2n}\right|^2 + b|h\mu|^2\,\mathrm{tr}\,\Phi_{2n}\Phi_{2n}$$

where $b = \frac{(\ln 4-1)}{8\pi^2}\widetilde{N}_c$ and we did not include Φ_{2n}-independent term and after differentiating this with respect to Φ_{2n} one obtains the local nonzero stable point given by

$$h\Phi_{2n} \simeq \frac{\mu_\phi}{b}\mathbf{1}_n\otimes i\sigma_2 \qquad \text{or} \qquad M_{2n}\simeq\frac{\alpha\Lambda^3}{\widetilde{N}_c}\mathbf{1}_n\otimes i\sigma_2$$

corresponding to the location of w coordinates for n flavor D4-branes, very close to the NS5-brane, between the $D6_{-\theta}$-branes and the NS5'-branes.

- Non-coincident $D6_{-\theta}$-branes and NS5'-branes

Let us consider when the numbers of $D6_{-\theta}$-branes and the number of NS5'-branes minus one are equal:$N_f = k'$. We displace the k' upper $D6_{-\theta}$-branes and upper NS5'-branes, given in Figure 4A, in the $+v$ direction respectively to two k' different points denoted by $v_{D6_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$(and their mirrors in the $-v$ direction). A single NS5'-brane is located at $v = 0$. Then there are $\sum_{j=1}^{k'} \widetilde{N}_{c,j} (\equiv N_{f,j} - N_{c,j} - 2) = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N_{f,j} = N_f$. When all the upper $NS5'_j$-branes and $D6_{-\theta,j}$-branes($j = 1, 2, \cdots, k'$) are distinct, the low energy physics corresponds to k' decoupled supersymmetric gauge theories with gauge groups $\prod_{j=1}^{k'} Sp(\widetilde{N}_{c,j})$.

One deforms the Figure 4B by displacing the multiple $D6_{-\theta}$-branes and $NS5'$-branes along v direction [15]. Then the n curved flavor D4-branes attached to them as well as other D4-branes are displaced also as k' different n_j's connecting between $D6_{-\theta,j}$-brane and $NS5'_j$-brane. Then the magnetic superpotential (3.7) can be rewritten in terms of Φ_j, q_j and $\phi'_{\widetilde{N}_{c,j}}$ which is coming from 4-4 strings connecting between the j-th $\widetilde{N}_{c,j}$ D4-branes

$$W_{dual} = \sum_{j=1}^{k'} \Big[h\Phi_j q_j q_j + \frac{\mu_\phi}{2} h^2 \, \mathrm{tr}\, \Phi_j^2 - h \, \mathrm{tr}\, \mu_j^2 \Phi_j + \frac{g_{\phi'}}{2} \, \mathrm{tr}\, {\phi'}_{\widetilde{N}_{c,j}}^{2k'+2} + q_j \phi'_{\widetilde{N}_{c,j}} q_j \Big] \tag{3.8}$$

with $\mu_j^2 = m_j \Lambda_j$ and $\mu_\phi = \alpha \Lambda^2$ as before.

One splits the $2(N_{f,j} - \widetilde{N}_{c,j} - l_j) \times 2(N_{f,j} - \widetilde{N}_{c,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $q_j q_j$ into blocks of size $2n_j$ and $2(N_{f,j} - \widetilde{N}_{c,j} - l_j - n_j)$ for all j as follows:

$$\begin{aligned} h\Phi \;&=\; \begin{pmatrix} 0_{2(\widetilde{N}_c+l+n)} & 0 & 0 & \cdots & 0 \\ 0 & \frac{\mu_1^2}{\mu_\phi} \mathbf{1}_{N_{f,1}-\widetilde{N}_{c,1}-l_1-n_1} \otimes i\sigma_2 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \frac{\mu_{k'}^2}{\mu_\phi} \mathbf{1}_{N_{f,k'}-\widetilde{N}_{c,k'}-l_{k'}-n_{k'}} \otimes i\sigma_2 \end{pmatrix} \\ &- \;\mathrm{diag}(\phi'_{\widetilde{N}_{c,1}}, \cdots, \phi'_{\widetilde{N}_{c,k'}}, 0_{N_f-\widetilde{N}_c}) \otimes i\sigma_2 \\ &+ \;\mathrm{diag}(0_{2(\widetilde{N}_c+l)}, h\Phi_{2n_1}, \cdots, h\Phi_{2n_{k'}}, 0_{2(N_f-\widetilde{N}_c-l-n)}) \end{aligned} \tag{3.9}$$

and

$$\begin{aligned} qq \;&=\; \begin{pmatrix} 0_{2\widetilde{N}_c} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \mu_1^2 \mathbf{1}_{2l_1} & 0 & 0 & \cdots & 0 & 0 \\ 0 & \cdot & \cdot & \cdot & \cdots & \cdot & 0 \\ 0 & 0 & 0 & 0 & \cdots & \mu_{k'}^2 \mathbf{1}_{2l_{k'}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0_{2(N_f-\widetilde{N}_c-l)} \end{pmatrix} \\ &+ \;\mathrm{diag}(\mu_1^2 \mathbf{1}_{\widetilde{N}_{c,1}} + \alpha\Lambda^2 \phi'_{\widetilde{N}_{c,1}}, \cdots, \mu_{k'}^2 \mathbf{1}_{\widetilde{N}_{c,k'}} + \alpha\Lambda^2 \phi'_{\widetilde{N}_{c,k'}}, 0_{N_f-\widetilde{N}_c}) \otimes i\sigma_2 \\ &+ \;\mathrm{diag}(0_{2(\widetilde{N}_c+l)}, \varphi_{2n_1}\varphi_{2n_1}, \cdots, \varphi_{2n_{k'}}\varphi_{2n_{k'}}, 0_{2(N_f-\widetilde{N}_c-l-n)}) \end{aligned} \tag{3.10}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$. Here φ_{2n_j} is $2n_j \times 2(\widetilde{N}_{c,j} - l_j)$ matrices and correspond to $2n_j$-flavors of fundamentals of the gauge group $Sp(\widetilde{N}_{c,j} - l_j)$ which is unbroken. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{2n_j} = \frac{\mu_j^2}{\mu_\phi} \mathbf{1}_{n_j} \otimes i\sigma_2$

and $\varphi_{2n_j} = 0$. The full one loop potential from (3.8), (3.9) and (3.10) can be written similarly and the local nonzero stable point arises as

$$h\Phi_{2n_j} \simeq \frac{\mu_\phi}{b_j}\mathbf{1}_{n_j} \otimes i\sigma_2 \qquad \text{or} \qquad M_{2n_j} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_{c,j}}\mathbf{1}_{n_j} \otimes i\sigma_2$$

corresponding to the w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane(and their mirrors).

Therefore, the meta-stable states, for fixed k' and θ, are classified by the number of various D4-branes and the positions of multiple $D6_{-\theta}$-branes and NS5'-branes: When N_f is not equal to k'(for example, when $N_f > k'$), then some coincident $D6_{-\theta}$-branes among N_f $D6_{-\theta}$-branes should be connected to the NS5'-branes in order to connect all the flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes.

4 $SU(N_c)$ with N_f-fund. and a symm.

4.1 Electric theory

The type IIA supersymmetric electric brane configuration [22, 23, 9] corresponding to $\mathcal{N} = 1$ $SU(N_c)$ gauge theory with a symmetric tensor S, a conjugate symmetric tensor $\widetilde{S}$, N_f-fundamental flavors $Q, \widetilde{Q}$ can be described as follows: one middle NS5-brane, $2k'$ NS5'-branes, N_c-color D4-branes, $2N_f$ D6-branes and the $O6^+$-plane(0123789). The N_c-color D4-branes are suspended between the left k' NS5'-branes and the right k' NS5'-branes and similarly each N_f D6-branes is located between the middle NS5-brane and the NS5'-branes.

Let us deform this theory [24] by adding the mass and the quartic terms for fundamental quarks. The former can be achieved by displacing the D6-branes along $\pm v$ direction leading to their coordinates $v = \pm v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by the angles $\mp\theta$ in (w, v)-plane and we denote them by $D6_{\mp\theta}$-branes. Therefore, in the electric gauge theory, the deformed superpotential is given by

$$W_{elec} = \frac{\alpha}{2}\,\mathrm{tr}(Q\widetilde{Q})^2 - m\,\mathrm{tr}\,Q\widetilde{Q} + \left[-\frac{\beta}{2}\,\mathrm{tr}(S\widetilde{S})^{k'+1} + m_S\,\mathrm{tr}\,S\widetilde{S}\right]. \tag{4.1}$$

Here the $\alpha = \frac{\tan\theta}{\Lambda}$ and $m = \frac{v_{D6_{-\theta}}}{2\pi\ell_s^2}$ correspond to the rotation and displacement of D6-branes and they are the same as the ones in [12]. The last two terms in (4.1) are due to the rotation of NS5'-branes where $\beta = \tan\omega$ and ω is a rotation angle of NS5'-branes in the (w, v)-plane and the relative displacement of D4-branes where $m_S = v_{NS5'}$ is the distance of D4-branes in v direction. When $k' = 1$ with small β limit and $m_S = 0$, this theory reduces to the one [12].

Therefore, we focus on the case with $k' \geq 2$ and $\beta, m_S \to 0$, i.e., we consider the multiple NS5'-branes with the electric superpotential (4.1) with this particular limit.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (4.1) in type IIA string theory as follows and draw this in Figure 5:

- One NS5-brane in (012345) directions with $w = 0 = x^6$
- $2k'$ NS5'-branes in (012389) directions with $v = 0$
- N_c-color D4-branes in (01236) directions with $v = 0 = w$
- N_f $D6_{\pm\theta}$-branes in (01237) directions and two other directions in (v, w)-plane
- $O6^+$-plane in (0123789) directions with $x^6 = 0 = v$



Figure 5: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c)$ with a symmetric tensor and fundamentals $Q, \widetilde{Q}$. Note that there are multiple k' NS5'-branes. A rotation of N_f D6-branes in (w, v)-plane, denoted by $D6_{\mp\theta}$-branes, corresponds to a quartic term for the fundamentals while a displacement of N_f D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals.

4.2 Magnetic theory

We move the $D6_{\mp\theta}$-branes and the left and right NS5'-branes through each other and use the linking numbers for the computation of creation of D4-branes as in [23]. Let us take the "extra" left- and right-$(k' - 1)$ NS5'-branes, compared with the single NS5'-brane case [23, 12], move them to the origin $x^6 = 0$, and rotate them by an angle $\frac{\pi}{2}$ in (w, v)-plane, coinciding with the middle NS5-brane. Temporarily, there exist $(2k' - 1)$ middle NS5-branes at $x^6 = 0$. Let us move the left $D6_\theta$-branes to the right all the way(and their mirrors, right $D6_{-\theta}$-branes to the left) past the NS5-branes and the right single NS5'-brane. Then the linking number l_m of a $D6_\theta$-brane becomes $l_m = \frac{1}{2} - n_{4L}$ while the one of the same $D6_\theta$-brane $l_e = -\frac{1}{2}$ in the electric theory. Then the number of D4-branes to the left of this $D6_\theta$-brane n_{4L} in the magnetic theory becomes 1 and we must add N_f D4-branes to the left side of all N_f $D6_\theta$-branes(and their mirrors). Note that at the $x^6 = 0$, the $D6_{\pm\theta}$-branes become the $D6_{\pm\frac{\pi}{2}}$-branes instantaneously [23, 12].

Then the extra $2(k'-1)$ middle NS5-branes, which were left- and right-$(k'-1)$ NS5'-branes in the electric theory, are moving to $\pm x^6$ direction by performing the remaining dual process and rotating by an angle $\frac{\pi}{2}$ in (w,v)-plane. This leads to the left k' NS5'-branes and the right k' NS5'-branes which look similar to the brane configuration of electric theory but we need to further take Seiberg dual for the remaining single NS5'-brane here. In this process there is a creation of D4-branes because the O6-plane, which has 4 D6-brane charge, is not parallel to these $2(k'-1)$ NS5-branes at $x^6=0$. Finally, after moving the left NS5'-brane, which does not participate in the dual process so far, to the right all the way past O6-plane(and its mirror, the right NS5'-brane to the left), the linking number of the NS5'-brane can be computed as $l_m = \frac{N_f}{2k'} + \frac{4(k'-1)}{2k'} - \frac{\widetilde{N}_c}{k'} + \frac{N_f}{k'}$. Note that the fractional contribution $\frac{k'-1}{k'}$ in the second term is due to the fact that only the left $(k'-1)$ NS5'-branes among k' NS5'-branes in the electric theory are crossing the O6-plane with an angle. Of course, the remaining left single NS5'-brane among k' NS5'-branes in the electric theory is crossing the N_f $D6_{-\theta}$-branes with an angle. Originally, the linking number was given by $l_e = -\frac{N_f}{2k'} - \frac{4(k'-1)}{2k'} + \frac{N_c}{k'}$. This leads to the fact that the number of D4-branes becomes $\widetilde{N}_c = 2N_f - N_c + 4(k'-1)$. Here the last constant term which depends on the number of NS5'-branes is due to the fact that the "extra" NS5'-branes are crossing the O6-plane. Of course, $k'=1$ case reduces to the known number of D4-branes $\widetilde{N}_c = 2N_f - N_c$ [23]. We present the Figure 6A.



Figure 6: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 5 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(6A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(6B). The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into (N_f-l)- and l- D4-branes(6A). Further n- D4-branes among (N_f-l)- D4-branes are moved to the NS5-brane(6B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c)$ gauge group and N_f-fundamental dual quarks $q,\widetilde{q}$ coming from 4-4 strings connecting between the color D4-branes and flavor D4-branes and a symmetric tensor and a conjugate symmetric tensor $s,\widetilde{s}$ coming from 4-4 strings connecting between the color D4-branes with $x^6<0$ and the color D4-branes with

$x^6 > 0$. Moreover, a single magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from 4-4 strings of flavor D4-branes. Then the magnetic superpotential with the limit $\beta, m_S \to 0$ is given by

$$W_{mag} = \frac{1}{\Lambda} M q(\widetilde{s}s)^{k'}\widetilde{q} + \frac{\alpha}{2}\,\mathrm{tr}\, M^2 - m\,\mathrm{tr}\, M. \tag{4.2}$$

The case where $k' = 1$ and $\alpha = 0$ leads to the previous result in [23]. Note that the number of color $\widetilde{N}_c$ depends on k'. In general, there are also different kinds of meson fields $M_j = Q(\widetilde{S}S)^j\widetilde{Q}, P_r = Q(\widetilde{S}S)^r\widetilde{S}Q$ and $\widetilde{P}_r = \widetilde{Q}S(\widetilde{S}S)^r\widetilde{Q}$ where $j = 1, 2, \cdots, k'$ and $r = 0, 1, \cdots, k'-1$ for general rotation angles of multiple NS5'-branes. The magnetic superpotential contains the interaction between these meson fields with $q, \widetilde{q}, s$ and $\widetilde{s}$ [24] as well as $(s\widetilde{s})^{k'+1}$ which will vanish for small β limit. However, the particular route we take above from an electric theory to the magnetic theory does not produce these meson fields because the N_f $D6_{\pm\theta}$-branes and the extra $2(k'-1)$ NS5'-branes do not create the D4-branes corresponding to those meson fields.

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (4.2) and the expectation values for M and $q\widetilde{s}s\widetilde{q}$ are obtained since this superpotential is the same as the one in [12]. Let us consider two different cases.

- Coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes(and their mirrors)

In this case, all the discussions given in [12] are satisfied with one exception that the number of colors here is different from the one in [12]. The theory has many nonsupersymmetric meta-stable ground states and when we rescale the meson field as $M = h\Lambda\Phi$, then the Kahler potential for Φ is canonical and the magnetic quarks are canonical near the origin of field space [1]. Then the magnetic superpotential can be written in terms of Φ or M

$$W_{mag} = h\Phi q(\widetilde{s}s)^{k'}\widetilde{q} + \frac{h^2\mu_\phi}{2}\,\mathrm{tr}\,\Phi^2 - h\mu^2\,\mathrm{tr}\,\Phi.$$

Now one splits the $(N_f - l) \times (N_f - l)$ block at the lower right corner of $h\Phi$ and $q\widetilde{s}s\widetilde{q}$ into blocks of size n and $(N_f - l - n)$ as follows [12]:

$$h\Phi = \begin{pmatrix} 0_l & 0 & 0 \\ 0 & h\Phi_n & 0 \\ 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{N_f-l-n} \end{pmatrix}, \qquad q(\widetilde{s}s)^{k'}\widetilde{q} = \begin{pmatrix} \mu^2\mathbf{1}_l & 0 & 0 \\ 0 & \varphi\widetilde{\beta}\beta\widetilde{\varphi} & 0 \\ 0 & 0 & 0_{N_f-l-n} \end{pmatrix}.$$

Here φ and $\widetilde{\varphi}$ are $n\times(\widetilde{N}_c - l)$ dimensional matrices and correspond to n flavors of fundamentals of the gauge group $SU(\widetilde{N}_c - l)$ which is unbroken. In the brane configuration shown in Figure 6B, they correspond to fundamental strings connecting the n flavor D4-branes and $(\widetilde{N}_c - l)$ color D4-branes [12]. The Φ_n and $\varphi\widetilde{\beta}\beta\widetilde{\varphi}$ are $n \times n$ matrices. The supersymmetric ground state corresponds to $h\Phi_n = \frac{\mu^2}{\mu_\phi}\mathbf{1}_n$ and $\varphi\widetilde{\beta} = 0 = \beta\widetilde{\varphi}$.

Now the full one loop potential for $\Phi_n, \hat{\varphi} \equiv \varphi\widetilde{\beta}$ and $\hat{\widetilde{\varphi}} \equiv \beta\widetilde{\varphi}$ [23] takes the form

$$\frac{V}{|h|^2} = |\Phi_n \hat{\varphi}|^2 + |\Phi_n \hat{\widetilde{\varphi}}|^2 + |\hat{\varphi}\hat{\widetilde{\varphi}} - \mu^2 \mathbf{1}_n + h\mu_\phi \Phi_n|^2 + b|h\mu|^2 \, \mathrm{tr}\, \Phi_n^\dagger \Phi_n,$$

where $b = \frac{(\ln 4 - 1)}{8\pi^2}\widetilde{N}_c$. Differentiating this potential with respect to $\Phi_n^\dagger$ and putting $\hat{\varphi} = 0 = \hat{\widetilde{\varphi}}$, one obtains

$$h\Phi_n \simeq \frac{\mu_\phi}{b}\mathbf{1}_n \qquad \text{or} \qquad M_n \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_c}\mathbf{1}_n \tag{4.3}$$

corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes(and their mirrors).

- Non-coincident $D6_{-\theta}$-branes and NS5'-branes

Let us consider the case where the numbers of $D6_{-\theta}$-branes and the NS5'-branes are equal to each other:$N_f = k'$. We displace the k' $D6_{-\theta}$-branes and NS5'-branes given in Figure 6A in the v direction respectively to two k' different points denoted by $v_{D6_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$, as in section 2. There are relations between the color and flavor D4-branes as follows: $\sum_{j=1}^{k'} \widetilde{N}_{c,j} = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N_{f,j} = N_f$.

One deforms the Figure 6B by displacing the multiple $D6_{-\theta}$-branes and $NS5'$-branes along v direction. Then the n curved flavor D4-branes attached to them are displaced also as k' different n_j's connecting between $D6_{-\theta,j}$-brane and $NS5'_j$-brane. Let us rescale the submeson field as $M_j = h\Lambda\Phi_j$ [15] and the Kahler potential for Φ_j is canonical and the magnetic quarks q_j and $\widetilde{q}_j$ as well as s_j and $\widetilde{s}_j$ are canonical near the origin of field space [1]. Then the magnetic superpotential can be rewritten in terms of $\Phi_j, q_j, \widetilde{q}_j, s_j$ and $\widetilde{s}_j$

$$W_{mag} = \sum_{j=1}^{k'} \left[h\Phi_j q_j (\widetilde{s}_j s_j)^{k'} \widetilde{q}_j + \frac{h^2 \mu_\phi}{2} \mathrm{tr}\, \Phi_j^2 - h\mu_j^2 \, \mathrm{tr}\, \Phi_j \right].$$

with $\mu_j^2 = m_j \Lambda_j$ and $\mu_\phi = \alpha\Lambda^2$ as before.

One splits the $(N_{f,j} - l_j) \times (N_{f,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $q_j \widetilde{s}_j s_j \widetilde{q}_j$ into blocks of size n_j and $(N_{f,j} - l_j - n_j)$ for all j as follows [15]:

$$\begin{aligned} h\Phi \;&=\; \begin{pmatrix} 0_{l+n} & 0 & 0 & \cdots & 0 \\ 0 & \frac{\mu_1^2}{\mu_\phi}\mathbf{1}_{N_{f,1}-l_1-n_1} & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \frac{\mu_{k'}^2}{\mu_\phi}\mathbf{1}_{N_{f,k'}-l_{k'}-n_{k'}} \end{pmatrix} \\ &+\; \mathrm{diag}(0_l, h\Phi_{n_1}, \cdots, h\Phi_{n_{k'}}, 0_{N_f-l-n}) \end{aligned}$$

and

$$\begin{aligned}
q(\widetilde{s}s)^{k'}\widetilde{q} &= \left(\begin{array}{cccccc}
\mu_1^2\mathbf{1}_{l_1} & 0 & 0 & \cdots & 0 & 0 \\
\cdot & \cdot & \cdot & \cdots & \cdot & 0 \\
0 & 0 & 0 & \cdots & \mu_{k'}^2\mathbf{1}_{l_{k'}} & 0 \\
0 & 0 & 0 & 0 & 0 & 0_{N_f-l}
\end{array}\right) \\
&+ \ \mathrm{diag}(0_l, \varphi_{n_1}\widetilde{\beta}_{n_1}\beta_{n_1}\widetilde{\varphi}_{n_1}, \cdots, \varphi_{n_{k'}}\widetilde{\beta}_{n'_k}\beta_{n'_k}\widetilde{\varphi}_{n_{k'}}, 0_{N_f-l-n})
\end{aligned}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$, as before. Here φ_{n_j} and $\widetilde{\varphi}_{n_j}$ are $n_j \times (\widetilde{N}_{c,j} - l_j)$ matrices and correspond to n_j-flavors of fundamentals of the gauge group $SU(\widetilde{N}_{c,j} - l_j)$ which is unbroken. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{n_j} = \frac{\mu_j^2}{\mu_\phi}\mathbf{1}_{n_j}$ and $\varphi_{n_j}\widetilde{\beta}_{n_j} = \beta_{n_j}\widetilde{\varphi}_{n_j} = 0$. The full one loop potential can be written similarly and the local nonzero stable point arises as

$$h\Phi_{n_j} \simeq \frac{\mu_\phi}{b_j}\mathbf{1}_{n_j} \qquad \text{or} \qquad M_{n_j} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_{c,j}}\mathbf{1}_{n_j}$$

corresponding to the nonzero w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane.

One can think of the brane configuration consisting of multiple outer NS5-branes as well as a single NS5'-brane, $O6^\pm$-planes and eight D6-branes. So we focus on the case where there is only a single outer NS5-brane [26, 25, 27, 28] when we discuss the other brane configurations containing this sub-brane configuration with NS5-brane, NS5'-brane, $O6^\pm$-planes and eight D6-branes later.

5 $SU(N_c) \times SU(N'_c)$ with N_f- and N'_f-fund. and bifund.

5.1 Electric theory

The type IIA supersymmetric electric brane configuration [29, 30, 31, 32, 33] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SU(N'_c)$ gauge theory with N_f-fundamental flavors $Q, \widetilde{Q}$, N'_f-fundamental flavors $Q', \widetilde{Q}'$ and bifundamentals $X, \widetilde{X}$ can be described as follows: one middle NS5-brane, $2k'$ NS5'-branes, N_c- and N'_c-D4-branes, and N_f- and N'_f-D6-branes. The N_c-color D4-branes are suspended between the middle NS5-brane and the right NS5'-branes, the N'_c-color D4-branes are suspended between the left NS5'-branes and the middle NS5-brane, the N_f D6-branes are located between the middle NS5-brane and the right NS5'-branes and the N'_f D6-branes are located between the left NS5'-branes and the middle NS5-brane.

Let us deform this theory by adding the mass term and the quartic term for fundamental quarks. The former can be achieved by displacing the D6-branes along $+v$ direction leading to their coordinates $v = +v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $D6_{-\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential is given by

$$\begin{aligned} W_{elec} &= \frac{\alpha}{2}\,\mathrm{tr}(Q\widetilde{Q})^2 - m\,\mathrm{tr}\,Q\widetilde{Q} + \frac{\alpha'}{2}\,\mathrm{tr}(Q'\widetilde{Q}')^2 - m'\,\mathrm{tr}\,Q'\widetilde{Q}' \\ &+ \left[-\frac{\beta}{2}\,\mathrm{tr}(X\widetilde{X})^{k'+1} + m_X\,\mathrm{tr}\,X\widetilde{X}\right]. \end{aligned} \tag{5.1}$$

The last two terms are due to the rotation of NS5'-branes where $\beta = (\tan\omega_L + \tan\omega_R)$ and the relative displacement of D4-branes where the mass $m_X = v_{NS5'}$ is the distance of D4-branes in v direction. When there are no D6-branes, this theory reduces to the one [15] and the case where there exists further restriction on the number of NS5'-branes $k' = 1$ has been studied in [14]. We focus on the case with $k' \geq 2$ and the limit $\beta, m_X \to 0$. When we take the Seiberg dual for the gauge group $SU(N_c)$, we put $\alpha' = 0$ and $m' = 0$(no displacement and no rotation of N'_f D6-branes) while for the Seiberg dual for the gauge group $SU(N'_c)$, we take $\alpha = 0$ and $m = 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (5.1) in type IIA string theory as follows and draw this in Figure 7:

- One middle NS5-brane in (012345) directions with $w = 0$
- $2k'$ NS5'-branes in (012389) directions $v = 0$
- N_f $D6_{-\theta}$-branes in (01237) directions and two other directions in (v, w)-plane
- N'_f D6-branes in (0123789) directions
- N_c- and N'_c-color D4-branes in (01236) directions with $v = 0 = w$



Figure 7: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SU(N'_c)$ with bifundamentals $X, \widetilde{X}$ and fundamentals $Q, \widetilde{Q}, Q', \widetilde{Q}'$. Note that there exist multiple $2k'$ NS5'-branes. A rotation of N_f D6-branes in (w, v)-plane corresponds to a quartic term for the fundamentals $Q, \widetilde{Q}$ while a displacement of N_f D6-branes in $+v$ direction corresponds to a mass term for the fundamentals $Q, \widetilde{Q}$.

5.2 Magnetic theory for $SU(N_c)$

After we move a middle NS5-brane to the right all the way past the right NS5'-branes, we arrive at the Figure 8A. Note that there exists a creation of N_f D4-branes connecting the N_f $D6_{-\theta}$-branes and the k' right NS5'-branes. The linking number of NS5-brane from Figure 8A is $l_m = \frac{N_f}{2} - \widetilde{N}_c$. On the other hand, the linking number of NS5-brane from Figure 7 is $l_e = -\frac{N_f}{2} + N_c - N'_c$. From these two relations, one obtains the number of colors of dual magnetic theory as follows [30]: $\widetilde{N}_c = N_f + N'_c - N_c$.



Figure 8: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration corresponding to Figure 7 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(8A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(8B) . The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into $(N_f - l)$- and l- D4-branes(8A). Further n- D4-branes among $(N_f - l)$- D4-branes are moved to the NS5-brane(8B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SU(N'_c)$ gauge group and N_f-fundamental dual quarks $q, \widetilde{q}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_f flavor D4-branes as well as $Q', \widetilde{Q}', Y$ and $\widetilde{Y}$ and gauge singlets. Moreover, a magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from 4-4 strings of N_f flavor D4-branes. Then the magnetic superpotential with the limit $\beta, m_X \to 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q \widetilde{q} + Y \widetilde{F}' \widetilde{q} + \widetilde{Y} q F' + \Phi' Y \widetilde{Y}\right] + \frac{\alpha}{2} \operatorname{tr} M^2 - mM. \tag{5.2}$$

The case with $k' = 1$ and $\alpha = 0$ was studied in [30]. Although the superpotential (5.2) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [30] appears as two things: nonzero α in (5.2) and multiple NS5'-branes in Figure 8. Here other meson fields are given by $\Phi' \equiv X\widetilde{X}, F' \equiv \widetilde{X}Q$ and $\widetilde{F}' \equiv X\widetilde{Q}$ [30].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (5.2) and the expectation values for M and $q\widetilde{q}$ are obtained. The F-term equations are almost the same as the one in [30] and the derivative of (5.2) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, F'$ and $\widetilde{F}'$ vanish as in [30].

- Coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

The theory has many nonsupersymmetric meta-stable ground states and when we rescale the meson field as $M = h\Lambda\Phi$ as before, then the Kahler potential for Φ is canonical and the magnetic quarks are canonical near the origin of field space [1]. Then the magnetic superpotential can be written as

$$W_{mag} = h\Phi q\widetilde{q} + \frac{h^2\mu_\phi}{2}\,\mathrm{tr}\,\Phi^2 - h\mu^2\,\mathrm{tr}\,\Phi + Y\widetilde{F}'\widetilde{q} + \widetilde{Y}qF' + \Phi'Y\widetilde{Y}$$

where $\mu^2 = m\Lambda$ and $\mu_\phi = \alpha\Lambda^2$. Now one splits the $(N_f - l) \times (N_f - l)$ block at the lower right corner of $h\Phi$ and $q\widetilde{q}$ into blocks of size n and $(N_f - l - n)$ as follows [7]:

$$h\Phi = \begin{pmatrix} 0_l & 0 & 0 \\ 0 & h\Phi_n & 0 \\ 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{N_f-l-n} \end{pmatrix}, \qquad q\widetilde{q} = \begin{pmatrix} \mu^2\mathbf{1}_l & 0 & 0 \\ 0 & \varphi\widetilde{\varphi} & 0 \\ 0 & 0 & 0_{N_f-l-n} \end{pmatrix}.$$

Here φ and $\widetilde{\varphi}$ are $n\times(\widetilde{N}_c-l)$ dimensional matrices and correspond to n flavors of fundamentals of the gauge group $SU(\widetilde{N}_c-l)$ which is unbroken. In the brane configuration shown in Figure 8B, they correspond to fundamental strings connecting the n flavor D4-branes and $(\widetilde{N}_c - l)$ color D4-branes [6]. The Φ_n and $\varphi\widetilde{\varphi}$ are $n \times n$ matrices. The supersymmetric ground state corresponds to $h\Phi_n = \frac{\mu^2}{\mu_\phi}\mathbf{1}_n$ and $\varphi = 0 = \widetilde{\varphi}$ [2].

Now the full one loop potential takes the form

$$\frac{V}{|h|^2} = |\Phi_n\varphi + Y\widetilde{F}'|^2 + |\Phi_n\widetilde{\varphi} + F'\widetilde{Y}|^2 + |\varphi\widetilde{\varphi} - \mu^2\mathbf{1}_n + h\mu_\phi\Phi_n|^2 + b|h\mu|^2\,\mathrm{tr}\,\Phi_n^\dagger\Phi_n,$$

where $b = \frac{(\ln 4-1)}{8\pi^2}\widetilde{N}_c$ and we do not write down Φ_n or $\Phi_n^\dagger$-independent terms. Differentiating this potential with respect to $\Phi_n^\dagger$ and putting $\varphi = 0 = \widetilde{\varphi}$, one obtains

$$h\Phi_n \simeq \frac{\mu_\phi}{b}\mathbf{1}_n \qquad \text{or} \qquad M_n \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_c}\mathbf{1}_n \tag{5.3}$$

corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes.

---

[2] Compared to a single gauge group, there exist extra k' left NS5'-branes, N_f' D6-branes and N_c' D4-branes. In Figure 8B, we consider the case where k' left NS5'-branes are far from k' right NS5'-branes. From the result of [34], in general, there exists a repulsive force between N_c' D4-branes and n D4-branes depending on their distance in w direction and an rotation angle between those D4-branes. In order to weaken this effect, we need to take the limit where their distance in w direction should be large and an rotation angle between them should be small. Furthermore, the distance beteen n D4-branes and the NS5-brane should be small also.

- Non-coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

When the numbers of $D6_{-\theta}$-branes and the NS5'-branes are equal to each other $N_f = k'$, we displace the k' $D6_{-\theta}$-branes and NS5'-branes given in Figure 8A in the v direction respectively to two k' different points denoted by $v_{D6_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$, as in section 2. There are $\sum_{j=1}^{k'} \widetilde{N}_{c,j} = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N_{f,j} = N_f$. Then the n curved flavor D4-branes attached to them are displaced also as k' different n_j's connecting between $D6_{-\theta,j}$-brane and $NS5'_j$-brane. Let us rescale the submeson field as $M_j = h\Lambda\Phi_j$ [15] and the Kahler potential for Φ_j is canonical and the magnetic quarks q_j and $\widetilde{q}_j$ are canonical near the origin of field space [1]. Then the magnetic superpotential can be rewritten in terms of Φ_j, q_j and $\widetilde{q}_j$

$$W_{mag} = \sum_{j=1}^{k'} \left[h\Phi_j q_j \widetilde{q}_j + \frac{h^2 \mu_\phi}{2} \operatorname{tr} \Phi_j^2 - h\mu_j^2 \operatorname{tr} \Phi_j \right] + \cdots$$

with $\mu_j^2 = m_j \Lambda_j$ and $\mu_\phi = \alpha\Lambda^2$ as before.

One splits the $(N_{f,j} - l_j) \times (N_{f,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $q_j\widetilde{q}_j$ into blocks of size n_j and $(N_{f,j} - l_j - n_j)$ for all j as follows [15]:

$$\begin{aligned} h\Phi &= \left(\begin{array}{ccccc} 0_{l+n} & 0 & 0 & \cdots & 0 \\ 0 & \frac{\mu_1^2}{\mu_\phi}\mathbf{1}_{N_{f,1}-l_1-n_1} & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \frac{\mu_{k'}^2}{\mu_\phi}\mathbf{1}_{N_{f,k'}-l_{k'}-n_{k'}} \end{array} \right) \\ &+ \ \operatorname{diag}(0_l, h\Phi_{n_1}, \cdots, h\Phi_{n_{k'}}, 0_{N_f-l-n}) \end{aligned}$$

and

$$\begin{aligned} q\widetilde{q} &= \left(\begin{array}{cccccc} \mu_1^2\mathbf{1}_{l_1} & 0 & 0 & \cdots & 0 & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot & 0 \\ 0 & 0 & 0 & \cdots & \mu_{k'}^2\mathbf{1}_{l_{k'}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0_{N_f-l} \end{array} \right) \\ &+ \ \operatorname{diag}(0_l, \varphi_{n_1}\widetilde{\varphi}_{n_1}, \cdots, \varphi_{n_{k'}}\widetilde{\varphi}_{n_{k'}}, 0_{N_f-l-n}) \end{aligned}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$. Here φ_{n_j} and $\widetilde{\varphi}_{n_j}$ are $n_j \times (\widetilde{N}_{c,j} - l_j)$ matrices and correspond to n_j-flavors of fundamentals of the gauge group $SU(\widetilde{N}_{c,j} - l_j)$ which is unbroken. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{n_j} = \frac{\mu_j^2}{\mu_\phi}\mathbf{1}_{n_j}$ and $\varphi_{n_j}\widetilde{\varphi}_{n_j} = 0$. The full one loop potential can be written similarly and the local nonzero stable point arises as

$$h\Phi_{n_j} \simeq \frac{\mu_\phi}{b_j}\mathbf{1}_{n_j} \qquad \text{or} \qquad M_{n_j} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_{c,j}}\mathbf{1}_{n_j} \tag{5.4}$$

corresponding to the nonzero w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane. Then, the meta-stable states, for fixed k' and θ, are specified by the number of various D4-branes and the positions of multiple $D6_{-\theta}$-branes and NS5'-branes.

6 $Sp(N_c) \times SO(2N'_c)$ with N_f-fund., N'_f-vectors. and bifund.

6.1 Electric theory

The type IIA supersymmetric electric brane configuration [35, 36, 37] corresponding to $\mathcal{N}=1$ $Sp(N_c) \times SO(2N'_c)$ gauge theory with N'_f-vectors Q', N_f-fundamental flavors Q and bifundamental X can be described as follows: one middle NS5-brane, $2(2k'+1)$ NS5'-branes, $2N_c$- and $2N'_c$-D4-branes, and $2N_f$- and $2N'_f$-D6-branes as well as $O4^{\pm}$-planes. The $2N_c$-color D4-branes are suspended between the middle NS5-brane and the right NS5'-branes, the $2N'_c$-color D4-branes are suspended between the left NS5'-branes and the middle NS5-brane, the $2N_f$ D6-branes are located between the middle NS5-brane and the right NS5'-branes and the $2N'_f$ D6-branes are located between the left NS5'-branes and the middle NS5-brane.

Let us deform this theory by adding the mass term and the quartic term for fundamental quarks. The former can be achieved by displacing the D6-branes along $\pm v$ direction leading to their coordinates $v = \pm v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $D6_{-\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential is given by

$$\begin{aligned} W_{elec} &= \frac{\alpha}{2} \operatorname{tr}(QQ)^2 - m \operatorname{tr} QQ + \frac{\alpha'}{2} \operatorname{tr}(Q'Q')^2 - m' \operatorname{tr} Q'Q' \\ &+ \left[-\frac{\beta}{2} \operatorname{tr}(XX)^{2k'+2} + m_X \operatorname{tr} XX \right]. \end{aligned} \tag{6.1}$$

The last two terms are due to the rotation of NS5'-branes where $\beta = \tan \omega$ and the relative displacement of D4-branes where the mass $m_X = v_{NS5'}$ is the distance of D4-branes in v direction. When there are no D6-branes, this theory reduces to the one [15] and the case where there exists a further restriction on the number of NS5'-branes $k' = 0$ has been studied in [14]. We focus on the case with $k' \geq 1$ and $\beta, m_X \to 0$. When we take the Seiberg dual for the gauge group $Sp(N_c)$, we put $\alpha' = 0$ and $m' = 0$ while for the Seiberg dual for the gauge group $SO(2N'_c)$, we take $\alpha = 0$ and $m = 0$ instead.

Let us summarize the $\mathcal{N}=1$ supersymmetric electric brane configuration with superpotential (6.1) in type IIA string theory as follows and draw this in Figure 9:

- One middle NS5-brane in (012345) directions with $w = 0$

- $2(2k'+1)$ NS5'-branes in (012389) directions $v=0$
- $2N_f$ $D6_{-\theta}$-branes in (01237) directions and two other directions in (v,w)-plane
- $2N'_f$ D6-branes in (0123789) directions
- $2N_c$- and $2N'_c$-color D4-branes in (01236) directions with $v=0=w$
- $O4^{\pm}$-planes in (01236) directions with $w=0=v$



Figure 9: The $\mathcal{N}=1$ supersymmetric electric brane configuration for the gauge group $Sp(N_c)\times SO(2N'_c)$ with bifundamental X and fundamentals and vectors Q, Q'. Note that there are multiple $2(2k'+1)$ NS5'-branes. A rotation of N_f D6-branes in (w,v)-plane corresponds to a quartic term for the fundamentals Q while a displacement of N_f D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals Q.

6.2 Magnetic theory for $Sp(N_c)$

After we move the middle NS5-brane to the right all the way past the right NS5'-branes, the linking number of NS5-brane from Figure 10A is given by $l_m = \frac{(2N_f)}{2} - 1 - (1) - 2\widetilde{N}_c$. Originally, it was given by $l_e = -\frac{(2N_f)}{2} + 1 - (-1) + 2N_c - 2N'_c$ from Figure 9. Therefore, by the linking number conservation and equating these two l's each other, we are left with the number of colors in the magnetic theory [35] $\widetilde{N}_c = N_f + N'_c - N_c - 2$.

The low energy theory on the color D4-branes has $Sp(\widetilde{N}_c)\times SO(2N'_c)$ gauge group and N_f-fundamental dual quarks q coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_f flavor D4-branes as well as Q', Y and gauge singlets. Moreover, a magnetic meson field $M \equiv QQ$ is $2N_f \times 2N_f$ matrix and comes from 4-4 strings of $2N_f$ flavor D4-branes. Then the magnetic superpotential with the limit $\beta, m_X \to 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} Mqq + Y\Phi' Y + qNY\right] + \frac{\alpha}{2}\,\mathrm{tr}\, M^2 - mM. \tag{6.2}$$

The case with $k'=0$ and $\alpha=0$ was studied in [35]. Although the superpotential (6.2) does not depend on the multiplicity $(2k'+1)$ of NS5'-branes in our particular limit, the difference from the previous result of [35] appears as 1) nonzero α in (6.2) and 2) multiple NS5'-branes in Figure 10A. Here other meson fields are given by $\Phi' \equiv XX$ and $N \equiv QX$.



Figure 10: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 9 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(10A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(10B) . The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into (N_f-l)- and l- D4-branes(10A). Further n- D4-branes among (N_f-l)- D4-branes are moved to the NS5-brane(10B).

One can compute the F-term equations for this superpotential (6.2) and the expectation values for M and qq are obtained. The F-term equations are almost the same as the one in [35] and the derivative of (6.2) with respect to the meson field M has α-dependent term. The vacuum expectation values for Y and N vanish as in [35].

- Coincident N_f $D6_{-\theta}$-branes and $(k'+\frac{1}{2})$ NS5'-branes

The theory has many nonsupersymmetric meta-stable ground states and when we rescale the meson field as $M=h\Lambda\Phi$ as before. Then the magnetic superpotential can be written as

$$W_{mag}=h\Phi qq+\frac{h^2\mu_\phi}{2}\operatorname{tr}\Phi^2-h\mu^2\operatorname{tr}\Phi+Y\Phi'Y+qNY$$

where $\mu^2=m\Lambda$ and $\mu_\phi=\alpha\Lambda^2$. Now one splits the $2(N_f-l)\times 2(N_f-l)$ block at the lower right corner of $h\Phi$ and qq into blocks of size $2n$ and $2(N_f-l-n)$ as follows [7]:

$$h\Phi=\left(\begin{array}{ccc} 0_{2l} & 0 & 0\\ 0 & h\Phi_{2n} & 0\\ 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{N_f-l-n}\otimes i\sigma_2\end{array}\right),\qquad qq=\left(\begin{array}{ccc} \mu^2\mathbf{1}_{2l} & 0 & 0\\ 0 & \varphi\varphi & 0\\ 0 & 0 & 0_{2(N_f-l-n)}\end{array}\right).$$

Here φ is $2n\times 2(\widetilde{N}_c-l)$ dimensional matrices and correspond to $2n$ flavors of fundamentals of the gauge group $Sp(\widetilde{N}_c-l)$ which is unbroken. The Φ_{2n} and $\varphi\varphi$ are $2n\times 2n$ matrices. The supersymmetric ground state corresponds to $h\Phi_{2n}=\frac{\mu^2}{\mu_\phi}\mathbf{1}_n\otimes i\sigma_2$ and $\varphi=0$.

Now the full one loop potential takes the form

$$\frac{V}{|h|^2}=|\Phi_{2n}\varphi+NY|^2+|\varphi\varphi-\mu^2\mathbf{1}_{2n}+h\mu_\phi\Phi_{2n}|^2+b|h\mu|^2\operatorname{tr}\Phi_{2n}\Phi_{2n},$$

where $b = \frac{(\ln 4 - 1)}{8\pi^2} \widetilde{N}_c$ and we do not write down Φ_{2n}-independent terms. Differentiating this potential with respect to Φ_{2n} and putting $\varphi = 0$, one obtains

$$h\Phi_{2n} \simeq \frac{\mu_\phi}{b} \mathbf{1}_n \otimes i\sigma_2 \qquad \text{or} \qquad M_{2n} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_c} \mathbf{1}_n \otimes i\sigma_2$$

corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes.

- Non-coincident N_f $D6_{-\theta}$-branes and $(k' + \frac{1}{2})$ NS5'-branes

When the numbers of $D6_{-\theta}$-branes and the number of NS5'-branes minus one are equal $N_f = k'$, we displace the k' upper $D6_{-\theta}$-branes and upper NS5'-branes, given in Figure 10A, in the $+v$ direction respectively to two k' different points denoted by $v_{D6_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$(and their mirrors in the $-v$ direction). A single NS5'-brane is located at $v = 0$. Then there exist $\sum_{j=1}^{k'} \widetilde{N}_{c,j} (\equiv N_{f,j} - N_{c,j} - 2) = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N_{f,j} = N_f$. When all the upper $NS5'_j$-branes and $D6_{-\theta,j}$-branes($j = 1, 2, \cdots, k'$) are distinct, the low energy physics corresponds to k' decoupled supersymmetric gauge theories with gauge groups $\left[\prod_{j=1}^{k'} Sp(\widetilde{N}_{c,j})\right] \times SO(2N'_c)$.

One deforms the Figure 10B by displacing the multiple $D6_{-\theta}$-branes and $NS5'$-branes along v direction [15]. Then the n curved flavor D4-branes attached to them are displaced also as k' different n_j's connecting between $D6_{-\theta,j}$-brane and $NS5'_j$-brane. Then the magnetic superpotential can be rewritten in terms of Φ_j, q_j

$$W_{mag} = \sum_{j=1}^{k'} \left[h\Phi_j q_j q_j + \frac{h^2 \mu_\phi}{2} \operatorname{tr} \Phi_j^2 - h\mu_j^2 \operatorname{tr} \Phi_j \right] + \cdots$$

with $\mu_j^2 = m_j \Lambda_j$ and $\mu_\phi = \alpha \Lambda^2$ as before.

One splits the $2(N_{f,j} - l_j) \times 2(N_{f,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $q_j q_j$ into blocks of size $2n_j$ and $2(N_{f,j} - l_j - n_j)$ for all j as follows:

$$\begin{aligned} h\Phi &= \begin{pmatrix} 0_{2(l+n)} & 0 & 0 & \cdots & 0 \\ 0 & \frac{\mu_1^2}{\mu_\phi} \mathbf{1}_{N_{f,1}-l_1-n_1} \otimes i\sigma_2 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \frac{\mu_{k'}^2}{\mu_\phi} \mathbf{1}_{N_{f,k'}-l_{k'}-n_{k'}} \otimes i\sigma_2 \end{pmatrix} \\ &+ \operatorname{diag}(0_{2l}, h\Phi_{2n_1}, \cdots, h\Phi_{2n_{k'}}, 0_{2(N_f-l-n)}) \end{aligned}$$

and

$$\begin{aligned} qq &= \begin{pmatrix} \mu_1^2 \mathbf{1}_{2l_1} & 0 & 0 & \cdots & 0 & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot & 0 \\ 0 & 0 & 0 & \cdots & \mu_{k'}^2 \mathbf{1}_{2l_{k'}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0_{2(N_f-l)} \end{pmatrix} \\ &+ \operatorname{diag}(0_{2l}, \varphi_{2n_1}\varphi_{2n_0}, \cdots, \varphi_{2n_{k'}}\varphi_{2n_{k'}}, 0_{2(N_f-l-n)}) \end{aligned}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$. Here φ_{2n_j} is $2n_j \times 2(\widetilde{N}_{c,j} - l_j)$ matrices and correspond to $2n_j$-flavors of fundamentals of the gauge group $Sp(\widetilde{N}_{c,j} - l_j)$ which is unbroken. Moreover, the Φ_{2n_j} and $\varphi_{2n_j}\varphi_{2n_j}$ are $2n_j \times 2n_j$ matrices. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{2n_j} = \frac{\mu_j^2}{\mu_\phi} \mathbf{1}_{n_j} \otimes i\sigma_2$ and $\varphi_{2n_j} = 0$. The full one loop potential can be written similarly and the local nonzero stable point arises as

$$h\Phi_{2n_j} \simeq \frac{\mu_\phi}{b_j} \mathbf{1}_{n_j} \otimes i\sigma_2 \qquad \text{or} \qquad M_{2n_j} \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_{c,j}} \mathbf{1}_{n_j} \otimes i\sigma_2$$

corresponding to the w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane(and their mirrors). Then, the meta-stable states, for fixed k' and θ, are specified by the number of various D4-branes and the positions of multiple $D6_{-\theta}$-branes and NS5'-branes.

7 $SU(N_c) \times SO(2N'_c)$ with N_f-fund., $2N'_f$-vectors and bi-fund.

7.1 Electric theory

The type IIA supersymmetric electric brane configuration [38, 30] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SO(2N'_c)$ gauge theory with N_f-fundamental flavors $Q, \widetilde{Q}$, $2N'_f$-vectors Q' and bifundamentals $X, \widetilde{X}$ can be described as follows: two NS5-branes, $2k'$ NS5'-branes, N_c- and $2N'_c$-D4-branes, $2N_f$- and $2N'_f$-D6-branes and $O6^+$-plane. The $2N'_c$-color D4-branes are suspended between the two NS5-branes and the N_c-color D4-branes are suspended between the NS5-brane and the NS5'-branes(and their mirrors).

Let us deform this theory by adding the mass term and the quartic term for the fundamentals. The former can be achieved by displacing the D6-branes along $+v$ direction leading to their coordinates $v = +v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $D6_{-\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential is given by

$$\begin{aligned} W_{elec} &= \frac{\alpha}{2} \operatorname{tr}(Q\widetilde{Q})^2 - m \operatorname{tr} Q\widetilde{Q} + \frac{\alpha'}{2} \operatorname{tr}(Q'Q')^2 - m' \operatorname{tr} Q'Q' \\ &+ \left[-\frac{\beta}{2} \operatorname{tr}(X\widetilde{X})^{k'+1} + m_X \operatorname{tr} X\widetilde{X} \right]. \end{aligned} \tag{7.1}$$

The last two terms are due to the rotation of NS5'-branes where $\beta = \tan\omega$ and the relative displacement of D4-branes where the mass $m_X = v_{NS5'}$ is the distance in v direction. This reduces to the one [30] when $k' = 1$ and $\alpha = 0 = \alpha' = m'$. We focus on the case with $k' \geq 2$

and $\beta, m_X \to 0$. When we take the Seiberg dual for the gauge group $SU(N_c)$, we put $\alpha' = 0$ and $m' = 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (7.1) in type IIA string theory as follows and draw this in Figure 11:

- Two NS5-branes in (012345) directions with $w = 0$
- $2k'$ NS5'-branes in (012389) directions $v = 0$
- N_f $D6_{\pm\theta}$-branes in (01237) directions and two other directions in (v, w)-plane
- $2N_f'$ D6-branes in (0123789) directions
- N_c- and $2N_c'$-color D4-branes in (01236) directions with $v = 0 = w$
- $O6^+$-plane in (0123789) directions with $x^6 = 0 = v$



Figure 11: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SO(2N_c')$ with bifundamental $X, \widetilde{X}$, fundamentals $Q, \widetilde{Q}$ and vectors Q'. Note that there are multiple k' NS5'-branes(and its mirrors). A rotation of N_f D6-branes in (w, v)-plane corresponds to a quartic term for the fundamentals while a displacement of N_f D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals.

7.2 Magnetic theory for $SU(N_c)$

After we move the right NS5-brane to the right all the way past the right NS5'-branes, we arrive at the Figure 12A. Note that there exists a creation of N_f D4-branes connecting the N_f $D6_{-\theta}$-branes and the k' right NS5'-branes. The linking number of NS5-brane from Figure 12A is $l_m = \frac{N_f}{2} - \widetilde{N}_c$. On the other hand, the linking number of NS5-brane from Figure 11 is $l_e = -\frac{N_f}{2} + N_c - N_c'$. From these two relations, one obtains the number of colors of dual magnetic theory as follows [30]: $\widetilde{N}_c = N_f + N_c' - N_c$.

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SO(2N_c')$ gauge group and N_f-fundamental dual quarks $q, \widetilde{q}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_f flavor D4-branes as well as $Q', Y, \widetilde{Y}$ and gauge singlets. Moreover, a magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from 4-4 strings of N_f flavor D4-branes.



Figure 12: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 11 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(12A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(12B) . The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into (N_f-l)- and l- D4-branes(12A). Further n- D4-branes among (N_f-l)- D4-branes are moved to the NS5-brane(12B).

Then the magnetic superpotential with the limit $\beta, m_X \rightarrow 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q\widetilde{q} + Q'\Phi'Q' + Y\widetilde{F}'\widetilde{q} + \widetilde{Y} q F' + \Phi' Y \widetilde{Y}\right] + \frac{\alpha}{2} \operatorname{tr} M^2 - mM. \tag{7.2}$$

The case with $k'=1$ and $\alpha=0$ was studied in [30]. Although the superpotential (7.2) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [30] appears as two things: nonzero α in (7.2) and multiple NS5'-branes in Figure 12. Here other meson fields are given by $\Phi' \equiv X\widetilde{X}, F' \equiv \widetilde{X}Q$ and $\widetilde{F}' \equiv X\widetilde{Q}$ [30].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (7.2) and the expectation values for M and $q\widetilde{q}$ are obtained. The F-term equations are almost the same as the one in [30] and the derivative of (7.2) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, F', \widetilde{F}'$ and Q' vanish as in [30].

- Coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

By following the description of subsection 5.2, one obtains the local nonzero stable point given by (5.3) corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes in Figure 12B.

- Non-coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

The local nonzero stable point arises as (5.4) corresponding to the nonzero w coordinates of n_j flavor D4-branes between the $D6_{-\theta,j}$-brane and the $NS5'_j$-brane.

8 $SU(N_c) \times SU(N'_c)$ with a symm. and bifund.

8.1 Electric theory

The type IIA supersymmetric electric brane configuration [39, 40, 41, 42] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SU(N'_c)$ gauge theory with a symmetric tensor S, a conjugate symmetric tensor $\widetilde{S}$ and bifundamentals $X, \widetilde{X}$ can be described as follows: $(2k+1)$ NS5-brane, $2k'$ NS5'-branes, N_c- and N'_c-D4-branes and $O6^+$-plane. The N_c-color D4-branes are suspended between the left NS5'-branes and the right NS5'-branes and the N'_c-color D4-branes are suspended between the right NS5'-branes and the right NS5-branes(and their mirrors).

Let us deform this theory by adding the mass term and the higher order term for the bifundamentals. The former can be achieved by displacing the NS5-branes along $+v$ direction leading to their coordinates $v = +v_{NS5}$ [9] while the latter can be obtained by rotating the NS5-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $NS5_{-\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential by including the mass term and higher order term for symmetric tensor matter is given by

$$W_{elec} = -\frac{\alpha}{2} \operatorname{tr}(X\widetilde{X})^{k+1} + m \operatorname{tr} X\widetilde{X} + \left[-\frac{\beta}{2} \operatorname{tr}(S\widetilde{S})^{k'+1} + m_S \operatorname{tr} S\widetilde{S} \right]. \tag{8.1}$$

The last two terms are due to the rotation of NS5'-branes where $\beta = \tan \omega$ and the relative displacement of D4-branes where the mass $m_S = v_{NS5'}$ is the distance in v direction. This reduces to the one [14] when $k = k' = 1$ and $\beta = m_S = 0$. We focus on the case with $k, k' \geq 2$ and $\beta, m_S \to 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (8.1) in type IIA string theory as follows and draw this in Figure 13:

- One NS5-brane in (012345) directions with $w = 0 = x^6$
- $2k'$ NS5'-branes in (012389) directions with $v = 0$
- k $NS5_{\pm\theta}$-branes in (012389) directions with $v = 0$
- N_c- and N'_c-color D4-branes in (01236) directions with $v = 0 = w$
- $O6^+$-plane in (0123789) directions with $x^6 = 0 = v$

8.2 Magnetic theory for $SU(N_c)$

We apply the Seiberg dual to the $SU(N_c)$ factor and the two k' NS5'-branes are interchanged each other. The linking number of the right NS5'-brane in Figure 14A is $l_m = -\frac{\widetilde{N}_c}{k'} + \frac{N'_c}{k'}$ and the linking number of the left NS5'-brane in Figure 13 is given by $l_e = \frac{N_c}{k'} - \frac{N'_c}{k'}$. This leads to the fact that the number of D4-branes becomes $\widetilde{N}_c = 2N'_c - N_c$.



Figure 13: The $\mathcal{N}=1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SU(N_c')$ with a symmetric tensor $S, \widetilde{S}$ and bifundamentals $X, \widetilde{X}$. Note that there are multiple k $NS5_{\pm\theta}$-branes and $2k'$ NS5'-branes. A rotation of k NS5-branes in (w, v)-plane corresponds to a higher term for the bifundamentals while a displacement of k NS5-branes in $\pm v$ direction corresponds to a mass term for the bifundamentals.



Figure 14: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 13 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(13A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(13B) . The N_c' flavor D4-branes connecting between $NS5_{-\theta}$-branes and NS5'-branes are splitting into $(N_c'-l)$- and l- D4-branes(14A). Further n- D4-branes among $(N_c'-l)$- D4-branes are moved to the NS5-brane(14B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SU(N_c')$ gauge group and N_c'-fundamental dual "quarks" $Y, \widetilde{Y}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_c' flavor D4-branes as well as s and $\widetilde{s}$. Moreover, a single magnetic meson field $M \equiv X\widetilde{X}$ is $N_c' \times N_c'$ matrix and comes from 4-4 strings of N_c' flavor D4-branes. Then the magnetic superpotential with the limit $\beta, m_S \to 0$ is given by

$$W_{dual} = \frac{1}{\Lambda} M Y \widetilde{Y} - \frac{\alpha}{2} M^{k+1} + mM. \tag{8.2}$$

The case with $k = 1 = k'$ and $\alpha = 0$ was studied in [39]. Although the superpotential (8.2) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [39] appears as two things: nonzero α in (8.2) which has explicit k dependent term and multiple $NS5_{-\theta}$-branes and NS5'-branes in Figure 14A. For the supersymmetric vacua, one can compute the F-term equations for this superpotential (8.2) and the expectation values for M and $Y\widetilde{Y}$ are obtained. The F-term equations are the same as the one in [15].

- Coincident k $NS5_{-\theta}$-branes and k' NS5'-branes

The theory has many nonsupersymmetric meta-stable ground states due to the fact that there exists an attractive gravitational interaction between the flavor D4-branes and the NS5-brane from the DBI action. When we rescale the meson field as $M = h\Lambda\Phi$, then the Kahler potential for Φ is canonical and the magnetic quarks Y and $\widetilde{Y}$ are canonical near the origin of field space. Then the magnetic superpotential (8.2) can be rewritten as

$$W_{dual} = h\Phi Y\widetilde{Y} + \frac{\mu_\phi}{2}h^{k+1}\,\mathrm{tr}\,\Phi^{k+1} - h\mu^2\,\mathrm{tr}\,\Phi \tag{8.3}$$

with the new couplings $\mu^2 = -m\Lambda$ and $\mu_\phi = -\alpha\Lambda^{k+1}$.

Now one splits the $(N_c' - l) \times (N_c' - l)$ block at the lower right corner of M and $Y\widetilde{Y}$ into blocks of size n and $(N_c' - l - n)$ and then they are rewritten as follows [7]:

$$h\Phi = \begin{pmatrix} 0\mathbf{1}_l & 0 & 0 \\ 0 & h\Phi_n & 0 \\ 0 & 0 & \left[\frac{2\mu^2}{\mu_\phi(k+1)}\right]^{\frac{1}{k}}\mathbf{1}_{N_c'-l-n} \end{pmatrix}, \qquad Y\widetilde{Y} = \begin{pmatrix} \mu^2\mathbf{1}_l & 0 & 0 \\ 0 & \varphi\widetilde{\varphi} & 0 \\ 0 & 0 & 0_{N_c'-l-n} \end{pmatrix}. \tag{8.4}$$

Here φ and $\widetilde{\varphi}$ are $n \times (\widetilde{N}_c - l)$ matrices and correspond to n-flavors of fundamentals of the gauge group $SU(\widetilde{N}_c - l)$ which is unbroken. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_n = \left[\frac{2\mu^2}{\mu_\phi(k+1)}\right]^{\frac{1}{k}}\mathbf{1}_n$ and $\varphi\widetilde{\varphi} = 0$.

The full one loop potential for Φ_n, φ, and $\widetilde{\varphi}$ from (8.3) and (8.4) including the one loop result [1] takes the form

$$\frac{V}{|h|^2} = |\Phi_n\varphi|^2 + |\Phi_n\widetilde{\varphi}|^2 + \left|\varphi\widetilde{\varphi} - \mu^2\mathbf{1}_n + \frac{(k+1)h^k}{2}\mu_\phi\Phi_n^k\right|^2 + b|h\mu|^2\,\mathrm{tr}\,\Phi_n^\dagger\Phi_n, \tag{8.5}$$

where the positive numerical constant b is given by $b = \frac{(\ln 4-1)}{8\pi^2}\widetilde{N}_c$. Differentiating this potential (8.5) with respect to $\Phi_n^\dagger$ and putting $\varphi = 0 = \widetilde{\varphi}$, one obtains

$$h\Phi_n \simeq \left[\frac{2b}{k(k+1)\mu_\phi}\right]^{\frac{1}{k-2}}\mathbf{1}_n \qquad \text{or} \qquad M_n \simeq \left[\frac{\widetilde{N}_c}{\alpha k(k+1)\Lambda^3}\right]^{\frac{1}{k-2}}\mathbf{1}_n. \tag{8.6}$$

It is evident that the $(N_c' - l - n)$ flavor D4-branes between the $NS5_{-\theta}$-branes and the NS5'-branes are related to the corresponding eigenvalues of $h\Phi$ (8.4), i.e., $\left[\frac{2\mu^2}{\mu_\phi(k+1)}\right]^{\frac{1}{k}}\mathbf{1}_{N_c'-l-n}$ and the intersection point between the $(N_c' - l - n)$ D4-branes and the NS5'-branes is also given by $(v, w) = (0, +v_{NS5_{-\theta}}\cot\theta)$.

- Non-coincident k $NS5_{-\theta}$-branes and k' NS5'-branes

When the numbers of $NS5_{-\theta}$-branes and the NS5'-branes are equal to each other $k = k'$, we displace the k' $NS5_{-\theta}$-branes and NS5'-branes given in Figure 14A in the v direction respectively to two k' different points denoted by $v_{NS5_{-\theta,j}}$ and $v_{NS5'_j}$ where $j = 1, 2, \cdots, k'$, as in section 2. There are $\sum_{j=1}^{k'} \widetilde{N}_{c,j} = \widetilde{N}_c$ and $\sum_{j=1}^{k'} N'_{c,j} = N'_c$.

One deforms the Figure 14B by displacing the multiple $NS5_{-\theta}$-branes and $NS5'$-branes along v direction. Then the n curved flavor D4-branes attached to them are displaced also as k' different n_j's connecting between $NS5_{-\theta,j}$-brane and $NS5'_j$-brane. Let us rescale the submeson field as $M_j = h\Lambda\Phi_j$ [15] and the Kahler potential for Φ_j is canonical and the magnetic quarks Y_j and $\widetilde{Y}_j$ are canonical near the origin of field space [1]. Then the magnetic superpotential can be rewritten in terms of Φ_j, Y_j and $\widetilde{Y}_j$

$$W_{mag} = \sum_{j=1}^{k} \left[h\Phi_j Y_j \widetilde{Y}_j + \frac{h^{k+1}\mu_\phi}{2} \operatorname{tr} \Phi_j^{k+1} - h\mu_j^2 \operatorname{tr} \Phi_j \right]$$

with $\mu_j^2 = -m_j\Lambda_j$ and $\mu_\phi = -\alpha\Lambda^{k+1}$ as before.

One splits the $(N'_{c,j} - l_j) \times (N'_{c,j} - l_j)$ block at the lower right corner of $h\Phi_j$ and $Y_j\widetilde{Y}_j$ into blocks of size n_j and $(N'_{c,j} - l_j - n_j)$ for all j as follows [15]:

$$\begin{aligned} h\Phi &= \left[\frac{2}{\mu_\phi(k+1)}\right]^{\frac{1}{k}} \begin{pmatrix} 0_{l+n} & 0 & 0 & \cdots & 0 \\ 0 & \mu_1^{\frac{2}{k}} \mathbf{1}_{N'_{c,1}-l_1-n_1} & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdots & 0 \\ 0 & 0 & 0 & 0 & \mu_k^{\frac{2}{k}} \mathbf{1}_{N'_{c,k}-l_k-n_k} \end{pmatrix} \\ &+ \operatorname{diag}(0_l, h\Phi_{n_1}, \cdots, h\Phi_{n_k}, 0_{N'_c-l-n}) \end{aligned}$$

and

$$\begin{aligned} Y\widetilde{Y} &= \begin{pmatrix} \mu_1^2 \mathbf{1}_{l_1} & 0 & 0 & \cdots & 0 & 0 \\ \cdot & \cdot & \cdot & \cdots & \cdot & 0 \\ 0 & 0 & 0 & \cdots & \mu_k^2 \mathbf{1}_{l_k} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0_{N'_c-l} \end{pmatrix} \\ &+ \operatorname{diag}(0_l, \varphi_{n_1}\widetilde{\varphi}_{n_1}, \cdots, \varphi_{n_k}\widetilde{\varphi}_{n_k}, 0_{N'_c-l-n}) \end{aligned}$$

where $l = \sum_{j=1}^{k'} l_j$ and $n = \sum_{j=1}^{k'} n_j$. Here φ_{n_j} and $\widetilde{\varphi}_{n_j}$ are $n_j \times (\widetilde{N}_{c,j} - l_j)$ matrices and correspond to n_j-flavors of fundamentals of the gauge group $SU(\widetilde{N}_{c,j} - l_j)$ which is unbroken. The supersymmetric ground state corresponds to the vacuum expectation values by $h\Phi_{n_j} = \left[\frac{2\mu_j^2}{\mu_\phi(k+1)}\right]^{\frac{1}{k}} \mathbf{1}_{n_j}$ and $\varphi_{n_j}\widetilde{\varphi}_{n_j} = 0$. The full one loop potential can be written similarly and the local nonzero stable point arises as

$$h\Phi_{n_j} \simeq \left[\frac{2b_j}{k(k+1)\mu_\phi}\right]^{\frac{1}{k-2}} \mathbf{1}_{n_j} \qquad \text{or} \qquad M_{n_j} \simeq \left[\frac{\widetilde{N}_{c,j}}{\alpha k(k+1)\Lambda_j^3}\right]^{\frac{1}{k-2}} \mathbf{1}_{n_j}$$

corresponding to the nonzero w coordinates of n_j flavor D4-branes between the $NS5_{-\theta,j}$-brane and the $NS5'_j$-brane. Then, the meta-stable states, for fixed k, k' and θ, are specified by the number of various D4-branes and the positions of multiple $NS5_{-\theta}$-branes and NS5'-branes.

9 $SU(N_c) \times SU(N'_c)$ with an antisymm., eight-fund. and bifund.

9.1 Electric theory

The type IIA supersymmetric electric brane configuration [39] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SU(N'_c)$ gauge theory with an antisymmetric tensor A, a conjugate symmetric tensor $\widetilde{S}$, eight fundamentals $\hat{Q}$ and bifundamentals $X, \widetilde{X}$ can be described as follows: Two NS5-branes, $(2k'+1)$ NS5'-branes, N_c- and N'_c-D4-branes, eight D6-branes and $O6^{\pm}$-planes. The N_c-color D4-branes are suspended between the left NS5-brane and the right NS5-brane and the N'_c-color D4-branes are suspended between the right NS5-brane and the right NS5'-branes(and their mirrors).

Let us deform this theory by adding the mass term and the higher order term for the bifundamentals. The former can be achieved by displacing the NS5'-branes along $+v$ direction leading to their coordinates $v = +v_{NS5'}$ [9] while the latter can be obtained by rotating the NS5'-branes [6] by an angle $-\theta$ in (w, v)-plane and we denote them by $NS5_{-\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential is given by

$$W_{elec} = -\frac{\alpha}{2} \operatorname{tr}(X\widetilde{X})^{k'+1} + m \operatorname{tr} X\widetilde{X} + \left[-\frac{\beta}{2} \operatorname{tr}(A\widetilde{S})^2 + \hat{Q}\widetilde{S}\hat{Q} \right]. \tag{9.1}$$

The third term is due to the rotation of NS5-branes where $\beta = \tan\omega$. This theory reduces to the one [14] when $k' = 1$ and $\beta = 0$. We focus on the case with $k' \geq 2$ and $\beta \to 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (9.1) in type IIA string theory as follows and draw this in Figure 15:

- Two NS5-branes in (012345) directions with $w = 0$
- One NS5'-brane in (012389) directions with $v = 0 = x^6$
- k' $NS5_{\pm\theta}$-branes in (012345) directions with $w = 0$
- N_c- and N'_c-color D4-branes in (01236) directions with $v = 0 = w$
- Eight half D6-branes in (0123789) directions with $x^6 = 0 = v$
- $O6^{\pm}$-planes in (0123789) directions with $x^6 = 0 = v$



Figure 15: The $\mathcal{N}=1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SU(N'_c)$ with an antisymmetric tensor $A, \widetilde{S}$, eight fundamentals $\hat{Q}$, and bifundamentals $X, \widetilde{X}$. Note that there are k' $NS5_{\pm\theta}$-branes. A rotation of k' NS5'-branes in (w, v)-plane corresponds to a higher term for the bifundamentals while a displacement of k' NS5-branes in $\pm v$ direction corresponds to a mass term for the bifundamentals.

9.2 Magnetic theory for $SU(N_c)$

We apply the Seiberg dual to the $SU(N_c)$ factor and the two NS5-branes are interchanged each other. The linking number of the right NS5-brane in Figure 16A is $l_m = \frac{4}{2} - \widetilde{N}_c + N'_c$ and the linking number of the left NS5-brane in Figure 15 is given by $l_e = -\frac{4}{2} + N_c - N'_c$. This leads to the fact that the number of D4-branes becomes [39] $\widetilde{N}_c = 2N'_c - N_c + 4$.



A

B

Figure 16: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 15 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(16A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(16B) . The N'_c flavor D4-branes connecting between $NS5_{-\theta}$-branes and NS5'-brane are splitting into $(N'_c - l)$- and l- D4-branes(16A). Further n- D4-branes among $(N'_c - l)$- D4-branes are moved to the NS5-brane(16B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SU(N'_c)$ gauge group and N'_c-fundamental dual "quarks" $Y, \widetilde{Y}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N'_c flavor D4-branes as well as $a, \widetilde{s}$ and $\hat{q}$. Moreover, a single magnetic meson field $M \equiv X\widetilde{X}$ is $N'_c \times N'_c$ matrix and comes from 4-4 strings of N'_c flavor D4-branes. Then the magnetic superpotential with the limit $\beta \rightarrow 0$ is given by

$$W_{dual} = \frac{1}{\Lambda} M Y \widetilde{Y} - \frac{\alpha}{2} M^{k'+1} + mM + \hat{q} \widetilde{s} \hat{q}. \tag{9.2}$$

The case with $k' = 1$ and $\alpha = 0$ was studied in [39]. The difference from the previous result of [39] appears as two things: nonzero α in (9.2) which has explicit k' dependent term and multiple $NS5_{\mp\theta}$-branes in Figure 16A. For the supersymmetric vacua, one can compute the F-term equations for this superpotential (9.2) and the expectation values for M and $Y\widetilde{Y}$ are obtained.

- Coincident k' $NS5_{-\theta}$-branes

One obtains the vacuum expectation value (8.6) with a replacement k by k' by applying the prescription of subsection 8.2. This provides the w coordinates of n flavor D4-branes between the $NS5_{-\theta}$-branes and the NS5'-brane in Figure 16B.

- Non-coincident k' $NS5_{-\theta}$-branes

These non-coincident $NS5_{-\theta}$-branes can be obtained by taking those quark masses being unequal. Then all the previous descriptions for the meta-stable states can be applied in this case also without any difficulty.

10 $SU(N_c) \times SU(N_c')$ with N_f- and N_f'-fund., a symm. and bifund.

10.1 Electric theory

The type IIA supersymmetric electric brane configuration [43] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SU(N_c')$ gauge theory with N_f-fundamental flavors $Q, \widetilde{Q}$, N_f'-fundamental flavors $Q', \widetilde{Q}'$, a symmetric tensor S, a conjugate symmetric tensor $\widetilde{S}$ and bifundamentals $X, \widetilde{X}$ can be described as follows: Three NS5-branes, $2k'$ NS5'-branes, N_c- and N_c'-D4-branes, and $2N_f$- and $2N_f'$-D6-branes and $O6^+$-plane. The N_c-color D4-branes are suspended between the left NS5'-branes and the right NS5'-branes, the N_c'-color D4-branes are suspended between the right NS5'-branes and the right NS5-brane(and their mirrors), the N_f D6-branes are located between the middle NS5-brane and the right NS5'-branes and the N_f' D6-branes are located between the right NS5'-branes and the right NS5-brane(and their mirrors).

Let us deform this theory by adding the mass term and the quartic term for fundamental quarks. The former can be achieved by displacing the D6-branes along $\pm v$ direction leading to their coordinates $v = \pm v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes [6] by an angle $\mp\theta$ in (w, v)-plane and we denote them by $D6_{\mp\theta}$-branes(or $D6_{\mp\theta'}$-branes for the second gauge group). Then, in the electric gauge theory, the deformed superpotential is given by

$$\begin{aligned} W_{elec} &= \frac{\alpha}{2} \operatorname{tr}(Q\widetilde{Q})^2 - m \operatorname{tr} Q\widetilde{Q} + \frac{\alpha'}{2} \operatorname{tr}(Q'\widetilde{Q}')^2 - m' \operatorname{tr} Q'\widetilde{Q}' \\ &+ \left[-\frac{\beta}{2} \operatorname{tr}(S\widetilde{S})^{k'+1} + m_S \operatorname{tr} S\widetilde{S} \right] + \left[-\frac{\gamma}{2} \operatorname{tr}(X\widetilde{X})^2 + m_X \operatorname{tr} X\widetilde{X} \right]. \end{aligned} \tag{10.1}$$

The last four terms are due to the rotation of NS5'-branes and NS5-branes where $\beta = \tan\omega'$ and $\gamma = \tan\omega$ and the relative displacement of D4-branes where the mass $m_S = v_{NS5'}$ and the mass $m_X = v_{NS5}$ are the distance in v direction. The case of $k' = 1$ and $\alpha = 0$ with $\beta = \gamma = m_S = m_X = 0$ was studied in [43]. We focus on the case with $k' \geq 2$ and $\beta, \gamma, m_S, m_X \to 0$. When we take the Seiberg dual for the gauge group $SU(N_c)$, we put $\alpha' = 0$ and $m' = 0$ and for the Seiberg dual on the gauge group $SU(N_c')$ we take $\alpha = 0$ and $m = 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (10.1) in type IIA string theory as follows and draw this in Figure 17:

- $2k'$ NS5'-branes in (012389) directions $v = 0$
- Three NS5-branes in (012345) directions $w = 0$
- N_f $D6_{\pm\theta}$-branes in (01237) directions and two other directions in (v, w)-plane
- N_f' $D6_{\pm\theta'}$-branes in (01237) directions and two other directions in (v, w)-plane
- N_c- and N_c'-color D4-branes in (01236) directions with $v = 0 = w$
- $O6^+$-plane in (0123789) directions with $x^6 = 0 = v$



Figure 17: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SU(N_c')$ with a symmetric tensor $S, \widetilde{S}$, bifundamentals $X, \widetilde{X}$ and fundamentals $Q, \widetilde{Q}, Q', \widetilde{Q}'$. Note that there are $2k'$ NS5'-branes. A rotation of $N_f(N_f')$ D6-branes in (w, v)-plane corresponds to a quartic term for the fundamentals $Q, \widetilde{Q}(Q', \widetilde{Q}')$ while a displacement of $N_f(N_f')$ D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals $Q, \widetilde{Q}(Q', \widetilde{Q}')$.

10.2 Magnetic theory for $SU(N_c)$

After the left NS5'-branes and $D6_\theta$-branes and the right NS5'-branes and $D6_{-\theta}$-branes are exchanged each other, we arrive at the Figure 18A. One reads off the number of colors of dual magnetic theory from section 4 by noticing the new N_c' dependence: $\widetilde{N}_c = 2(N_f + N_c') - N_c + 4(k' - 1)$.



Figure 18: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration corresponding to Figure 17 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(18A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(18B) . The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-branes are splitting into $(N_f - l)$- and l- D4-branes(18A). Further n- D4-branes among $(N_f - l)$- D4-branes are moved to the NS5-brane(18B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SU(N_c')$ gauge group and N_f-fundamental dual quarks $q, \tilde{q}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_f flavor D4-branes as well as $Q', \widetilde{Q}', s, \tilde{s}, Y$ and $\widetilde{Y}$ and gauge singlets. Moreover, a magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from 4-4 strings of N_f flavor D4-branes. Then the magnetic superpotential with the limit $\beta, \gamma, m_S, m_X \to 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q\tilde{s}s\tilde{q} + Y\widetilde{F}'\tilde{q} + \widetilde{Y}qF' + \Phi' Y\widetilde{Y}\right] + \frac{\alpha}{2}\,\mathrm{tr}\, M^2 - mM. \tag{10.2}$$

The case with $k' = 1$ and $\alpha = 0$ was studied in [43] as mentioned before. Although the superpotential (10.2) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [43] appears as 1) nonzero α in (10.2) and 2) mutiple NS5'-branes in Figure 18A. Here other meson fields are given by $\Phi' \equiv X\widetilde{X}, F' \equiv \widetilde{X}Q$ and $\widetilde{F}' \equiv X\widetilde{Q}$ [43].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (10.2) and the expectation values for M and $q\tilde{s}s\tilde{q}$ are obtained. The F-term equations are almost the same as the one in [43] and the derivative of (10.2) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, F'$ and $\widetilde{F}'$ vanish as in [43].

- Coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

One gets the local stable point (4.3) corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes in Figure 18B by following the prescription of subsection 4.2.

- Non-coincident N_f $D6_{-\theta}$-branes and k' NS5'-branes

The local nonzero stable point arises similarly.

10.3 Magnetic theory for $SU(N_c')$

After we move the right NS5'-branes to the right all the way past the right NS5-brane(and its mirrors to the left), we arrive at the Figure 19A. Note that there exists a creation of $k'N_f'$ D4-branes connecting the N_f' $D6_{-\theta'}$-branes and the NS5'-brane. The linking number of NS5'-brane from Figure 19A is $l_m = \frac{N_f'}{2} - \frac{\widetilde{N}_c}{k'}$. On the other hand, the linking number of NS5'-brane from Figure 17 is $l_e = -\frac{N_f'}{2} + \frac{N_c'}{k'} - \frac{N_c}{k'}$. From these two relations, one obtains the number of colors of dual magnetic theory $\widetilde{N}_c' = k'N_f' + N_c - N_c'$. In this subsection we consider only the single number of NS5'-brane $k' = 1$ in order to deal with a single meson field.



Figure 19: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration corresponding to Figure 18 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(19A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(19B). The N_f' flavor D4-branes connecting between $D6_{-\theta'}$-branes and NS5'-brane are splitting into $(N_f'-l)$- and l- D4-branes(19A). Further n- D4-branes among $(N_f' - l)$- D4-branes are moved to the NS5-brane(19B).

The low energy theory on the color D4-branes has $SU(N_c) \times SU(\widetilde{N}_c')$ gauge group and N_f'-fundamental dual quarks $q', \widetilde{q}'$ coming from 4-4 strings connecting between the color $\widetilde{N}_c'$ D4-branes and N_f' flavor D4-branes as well as $Q, \widetilde{Q}, S, \widetilde{S}, Y, \widetilde{Y}$ and gauge singlets. Moreover, a magnetic meson field $M \equiv Q'\widetilde{Q}'$ is $N_f' \times N_f'$ matrix and comes from 4-4 strings of N_f' flavor D4-branes. Then the magnetic superpotential with the limit $\beta, \gamma, m_S, m_X \to 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q' \widetilde{q}' + Y F q' + \widetilde{Y} \widetilde{q}' \widetilde{F} + \Phi' Y \widetilde{Y}\right] + \frac{\alpha}{2} \operatorname{tr} M^2 - m' M. \tag{10.3}$$

The case with $k' = 1$ and $\alpha = 0$ was studied in [43]. Although the superpotential (10.3) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [43] appears as nonzero α in (10.3). Here other meson fields are given by $\Phi' \equiv X\widetilde{X}, F \equiv XQ'$ and $\widetilde{F} \equiv \widetilde{X}\widetilde{Q}'$ [43].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (10.3) and the expectation values for M and $q'\widetilde{q}'$ are obtained. The F-term equations are almost the same as the one in [43] and the derivative of (10.3) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, F$ and $\widetilde{F}$ vanish as in [43].

- Coincident N_f' $D6_{-\theta'}$-branes

One obtains the local stable point as (5.3) by changing the role of Q and $\widetilde{Q}$ into the one of Q' and $\widetilde{Q}'$. This gives the w coordinates of n flavor D4-branes between the $D6_{-\theta'}$-branes and the NS5'-brane in Figure 19B.

- Non-coincident N_f' $D6_{-\theta'}$-branes

These non-coincident $D6_{-\theta}$-branes can be obtained by taking those quark masses being unequal. Then all the previous descriptions for the meta-stable states can be applied in this case also without any difficulty.

11 $SU(N_c) \times SU(N_c')$ with N_f- and N_f'-fund., an antisymm., eight-fund. and bifund.

11.1 Electric theory

The type IIA supersymmetric electric brane configuration [43] corresponding to $\mathcal{N} = 1$ $SU(N_c) \times SU(N_c')$ gauge theory with N_f-fundamental flavors $Q, \widetilde{Q}$, N_f'-fundamental flavors $Q', \widetilde{Q}'$, an antisymmetric tensor A, a conjugate symmetric tensor $\widetilde{S}$, eight fundamentals $\hat{Q}$ and bifundamentals $X, \widetilde{X}$ can be described as follows: Two NS5-branes, $(2k' + 1)$ NS5'-branes, N_c- and N_c'-D4-branes, and $2N_f$- and $2N_f'$-D6-branes, eight D6-branes and $O6^{\pm}$-planes. The N_c-color D4-branes are suspended between the two NS5-branes, the N_c'-color D4-branes are suspended between the right NS5-brane and the right NS5'-branes(and their mirrors), the N_f D6-branes are located between the middle NS5'-brane and the right NS5-brane and the N_f' D6-branes are located between the right NS5-brane and the right NS5'-branes(and their mirrors).

Let us deform this theory by adding the mass term and the quartic term for fundamental quarks. The former can be achieved by displacing the D6-branes along $\pm v$ direction leading to their coordinates $v = \pm v_{D6}$ [9] while the latter can be obtained by rotating the D6-branes

[6] by an angle $\mp\theta$ in (w,v)-plane and we denote them by $D6_{\mp\theta}$-branes. Then, in the electric gauge theory, the general deformed superpotential is given by

$$\begin{aligned} W_{elec} &= \frac{\alpha}{2}\,\mathrm{tr}(Q\widetilde{Q})^2 - m\,\mathrm{tr}\,Q\widetilde{Q} + \hat{Q}\widetilde{S}\hat{Q} + \frac{\alpha'}{2}\,\mathrm{tr}(Q'\widetilde{Q}')^2 - m'\,\mathrm{tr}\,Q'\widetilde{Q}' \\ &- \frac{\beta}{2}\,\mathrm{tr}(A\widetilde{S})^2 + \left[-\frac{\gamma}{2}\,\mathrm{tr}(X\widetilde{X})^{k'+1} + m_X\,\mathrm{tr}\,X\widetilde{X}\right]. \end{aligned} \tag{11.1}$$

The last three terms are due to the rotation of NS5-branes and NS5'-branes where $\beta = \tan\omega$ and $\gamma = \tan\omega'$ and the relative displacement of D4-branes where the mass $m_X = v_{NS5'}$ is the distance in v direction. The case of $k' = 1$ and $\alpha = 0$ with $\beta = \gamma = m_X = 0$ was studied in [43]. We focus on the case with $k' \geq 2$ and $\beta, \gamma, m_X \to 0$. When we take the Seiberg dual for the gauge group $SU(N_c)$, we put $\alpha' = 0$ and $m' = 0$ and for the Seiberg dual on the gauge group $SU(N_c')$ we take $\alpha = 0$ and $m = 0$.

Let us summarize the $\mathcal{N} = 1$ supersymmetric electric brane configuration with superpotential (11.1) in type IIA string theory as follows and draw this in Figure 20:

- Two NS5-branes in (012345) directions with $w = 0$
- $(2k'+1)$ NS5'-branes in (012389) directions $v = 0$
- N_f $D6_{\pm\theta}$-branes in (01237) directions and two other directions in (v,w)-plane
- N_f' $D6_{\pm\theta'}$-branes in (01237) directions and two other directions in (v,w)-plane
- Eight D6-branes in (0123789) directions
- N_c- and N_c'-color D4-branes in (01236) directions with $v = 0 = w$
- $O6^{\pm}$-planes in (0123789) directions with $x^6 = 0 = v$



Figure 20: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the gauge group $SU(N_c) \times SU(N_c')$ with an antisymmetric tensor $A, \widetilde{S}$, bifundamentals $X, \widetilde{X}$, fundamentals $Q, \widetilde{Q}, Q', \widetilde{Q}'$ and fundamentals $\hat{Q}$. Note that there are multiple $2k'$ $NS5'$-branes. A rotation of N_f D6-branes in (w,v)-plane corresponds to a quartic term for the fundamentals $Q, \widetilde{Q}$ while a displacement of N_f D6-branes in $\pm v$ direction corresponds to a mass term for the fundamentals $Q, \widetilde{Q}$.

11.2 Magnetic theory for $SU(N_c)$

After the left NS5-branes and $D6_\theta$-branes and the right NS5-branes and $D6_{-\theta}$-branes are exchanged each other, we arrive at the Figure 21A. The number of colors of dual magnetic theory is given by [43] $\widetilde{N}_c = 2(N_f + N_c') - N_c + 4$.



Figure 21: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration corresponding to Figure 20 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(21A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(21B). The N_f flavor D4-branes connecting between $D6_{-\theta}$-branes and NS5'-brane are splitting into $(N_f - l)$- and l- D4-branes(21A). Further n-D4-branes among $(N_f - l)$- D4-branes are moved to the NS5-brane(22B).

The low energy theory on the color D4-branes has $SU(\widetilde{N}_c) \times SU(N_c')$ gauge group and N_f-fundamental dual quarks $q, \widetilde{q}$ coming from 4-4 strings connecting between the color $\widetilde{N}_c$ D4-branes and N_f flavor D4-branes as well as $Q', \widetilde{Q}', a, \widetilde{s}, \hat{q}, Y, \widetilde{Y}$ and gauge singlets. Moreover, a magnetic meson field $M \equiv Q\widetilde{Q}$ is $N_f \times N_f$ matrix and comes from 4-4 strings of N_f flavor D4-branes. Then the magnetic superpotential with the limit $\beta, \gamma, m_X \to 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q \widetilde{s} a \widetilde{q} + \hat{q} \widetilde{s} \hat{q} + Y \widetilde{F}' \widetilde{q} + \widetilde{Y} q F' + \Phi' Y \widetilde{Y} + \widetilde{M} \hat{q} \widetilde{q} \right] + \frac{\alpha}{2} \operatorname{tr} M^2 - mM. \qquad (11.2)$$

The case with $k' = 1$ and $\alpha = 0$ was studied in [43] as mentioned before. Although the superpotential (11.2) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result appears as 1) nonzero α in (11.2) and 2) multiple NS5'-branes in Figure 21A. Here other meson fields are given by $\Phi' \equiv X\widetilde{X}$, $F' \equiv \widetilde{X}Q$, $\widetilde{M} \equiv \hat{Q}\widetilde{Q}$ and $\widetilde{F}' \equiv X\widetilde{Q}$ [43].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (11.2) and the expectation values for M and $q\widetilde{s}a\widetilde{q}$ are obtained. The F-term equations are almost the same as the one in [43] and the derivative of (11.2) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, \hat{q}, \widetilde{M}, F'$ and $\widetilde{F}'$ vanish as in [43].

- Coincident N_f $D6_{-\theta}$-branes

The theory has many nonsupersymmetric meta-stable ground states and when we rescale the meson field as $M = h\Lambda\Phi$ as before, then the Kahler potential for Φ is canonical and the magnetic quarks are canonical near the origin of field space [1]. Then the magnetic superpotential can be written as

$$W_{mag} = h\Phi q\widetilde{s}a\widetilde{q} + \frac{h^2\mu_\phi}{2}\,\mathrm{tr}\,\Phi^2 - h\mu^2\,\mathrm{tr}\,\Phi + \hat{q}\widetilde{s}\hat{q} + YFq' + \widetilde{Y}\widetilde{q}'\widetilde{F}' + \Phi' Y\widetilde{Y} + \widetilde{M}\hat{q}\widetilde{q}$$

where $\mu^2 = m\Lambda$ and $\mu_\phi = \alpha\Lambda^2$. Now one splits the $(N_f - l) \times (N_f - l)$ block at the lower right corner of $h\Phi$ and $q\widetilde{s}a\widetilde{q}$ into blocks of size n and $(N_f - l - n)$ as follows [7]:

$$h\Phi = \begin{pmatrix} 0_l & 0 & 0 \\ 0 & h\Phi_n & 0 \\ 0 & 0 & \frac{\mu^2}{\mu_\phi}\mathbf{1}_{N_f-l-n} \end{pmatrix}, \qquad q\widetilde{s}a\widetilde{q} = \begin{pmatrix} \mu^2\mathbf{1}_l & 0 & 0 \\ 0 & \varphi\widetilde{\beta}\gamma\widetilde{\varphi} & 0 \\ 0 & 0 & 0_{N_f-l-n} \end{pmatrix}.$$

Here φ and $\widetilde{\varphi}$ are $n\times(\widetilde{N}_c - l)$ dimensional matrices and correspond to n flavors of fundamentals of the gauge group $SU(\widetilde{N}_c - l)$ which is unbroken. The Φ_n and $\varphi\widetilde{\beta}\gamma\widetilde{\varphi}$ are $n \times n$ matrices. The supersymmetric ground state corresponds to $h\Phi_n = \frac{\mu^2}{\mu_\phi}\mathbf{1}_n$ and $\varphi\widetilde{\beta} = 0 = \gamma\widetilde{\varphi}$.

Now the full one loop potential takes the form

$$\frac{V}{|h|^2} = |\Phi_n\varphi\widetilde{\beta} + \widetilde{Y}\widetilde{F}|^2 + |\Phi_n\gamma\widetilde{\varphi} + FY|^2 + |\varphi\widetilde{\beta}\gamma\widetilde{\varphi} - \mu^2\mathbf{1}_n + h\mu_\phi\Phi_n|^2 + b|h\mu|^2\,\mathrm{tr}\,\Phi_n^\dagger\Phi_n,$$

where $b = \frac{(\ln 4 - 1)}{8\pi^2}\widetilde{N}_c$ and we do not write down Φ_n or $\Phi_n^\dagger$-independent terms. Differentiating this potential with respect to $\Phi_n^\dagger$ and putting $\varphi\widetilde{\beta} = 0 = \gamma\widetilde{\varphi}$, one obtains

$$h\Phi_n \simeq \frac{\mu_\phi}{b}\mathbf{1}_n \qquad \text{or} \qquad M_n \simeq \frac{\alpha\Lambda^3}{\widetilde{N}_c}\mathbf{1}_n$$

corresponding to the w coordinates of n curved flavor D4-branes between the $D6_{-\theta}$-branes and the NS5'-branes.

- Non-coincident N_f $D6_{-\theta}$-branes

These non-coincident $D6_{-\theta}$-branes can be obtained by taking those quark masses being unequal.

11.3 Magnetic theory for $SU(N_c')$

After we move the right NS5-brane to the right all the way past the right NS5'-branes(and its mirrors to the left), we arrive at the Figure 22A. The linking number of NS5-brane from Figure 22A is $l_m = \frac{N_f'}{2} - \widetilde{N}_c$. On the other hand, the linking number of NS5-brane from Figure



Figure 22: The $\mathcal{N}=1$ supersymmetric magnetic brane configuration corresponding to Figure 20 with a misalignment between D4-branes when the gravitational potential of the NS5-brane is ignored(22A) and nonsupersymmetric brane configuration when the gravitational potential of the NS5-brane is considered(22B). Note that there are multiple $2k'$ $NS5'$-branes. The N_f' flavor D4-branes connecting between $D6_{-\theta'}$-branes and NS5'-brane are splitting into $(N_f'-l)$- and l- D4-branes(22A). Further n- D4-branes among $(N_f'-l)$- D4-branes are moved to the NS5-brane(22B).

21 is $l_e = -\frac{N_f'}{2} + N_c - N_c'$. From these two relations, one obtains the number of colors of dual magnetic theory $\widetilde{N}_c' = N_f' + N_c - N_c'$.

The low energy theory on the color D4-branes has $SU(N_c) \times SU(\widetilde{N}_c')$ gauge group and N_f'-fundamental dual quarks $q', \widetilde{q}'$ coming from 4-4 strings connecting between the color $\widetilde{N}_c'$ D4-branes and N_f' flavor D4-branes as well as $Q, \widetilde{Q}, A, \widetilde{S}, \hat{Q}, Y, \widetilde{Y}$ and gauge singlets. Moreover, a single magnetic meson field $M \equiv Q'\widetilde{Q}'$ is $N_f' \times N_f'$ matrix and comes from 4-4 strings of N_f' flavor D4-branes. Then the magnetic superpotential with the limit $\beta, \gamma, m_X \rightarrow 0$ is given by

$$W_{dual} = \left[\frac{1}{\Lambda} M q' \widetilde{q}' + Y F q' + \widetilde{Y} \widetilde{q}' \widetilde{F} + \Phi' Y \widetilde{Y} \right] + \frac{\alpha}{2} \operatorname{tr} M^2 - m' M. \tag{11.3}$$

The case with $k'=1$ and $\alpha=0$ was studied in [43]. Although the superpotential (11.3) does not depend on the multiplicity k' of NS5'-branes in our particular limit, the difference from the previous result of [43] appears as two things: nonzero α in (11.3) and multiple NS5'-branes in Figure 22A. Here other meson fields are given by $\Phi' \equiv X\widetilde{X}, F \equiv XQ'$ and $\widetilde{F} \equiv \widetilde{X}\widetilde{Q}'$ [43].

For the supersymmetric vacua, one can compute the F-term equations for this superpotential (11.3) and the expectation values for M and $q'\widetilde{q}'$ are obtained. The F-term equations are almost the same as the one in [43] and the derivative of (11.3) with respect to the meson field M has α dependent term. The vacuum expectation values for $Y, \widetilde{Y}, F$ and $\widetilde{F}$ vanish as in [43].

- Coincident N_f' $D6_{-\theta'}$-branes

One obtains the stable point (5.3) with appropriate replacement of N_f $D6_{-\theta}$-branes by the corresponding N_f' $D6_{-\theta'}$-branes. This gives the w coordinates of n curved flavor D4-branes between the $D6_{-\theta'}$-branes and the NS5'-branes in Figure 22B.

• Non-coincident N_f' $D6_{-\theta'}$-branes and k' NS5'-branes

The local nonzero stable point arises as (5.4).

12 Conclusions and outlook

As mentioned in the abstract, let us summarize the new features we have obtained from the meta-stable brane configurations.

Compared with the ones of [7, 6], the Figure 2 contains the multiple k' NS5'-branes and the presence of $\widetilde{N}_c$ flavor D4-branes connecting $D6_{-\theta}$-branes and NS5'-branes. Similarly, its symplectic version, characterized by the Figure 4, contains the multiple $(2k'+1)$ NS5'-branes and the presence of $\widetilde{N}_c$ flavor D4-branes connecting $D6_{-\theta}$-branes and NS5'-branes(and its mirrors). Compared with the one of [12], the Figure 6 has the multiple k' NS5'-branes and the different value of $\widetilde{N}_c$ which has k'-dependence. The Figures 8, 10, 12 have multiple NS5'-branes and rotated D6-branes, compared with [30].

Compared with [39], the Figure 14 has multiple rotated NS5-branes and multiple NS5'-branes while the Figure 16 has multiple rotated NS5'-branes. The Figure 18 contains the multiple k' NS5'-branes and the different value of $\widetilde{N}_c$ which has k'-dependence while the Figure 19 has only rotated D6-branes, compared with the previous result in [43]. Finally, the Figure 21 has only rotated D6-branes and the Figure 22 multiple NS5'-branes and rotated D6-branes, compared with [43].

We make some comments for the future directions along the line of meta-stable brane construction.

• As we mentioned, the construction for possible multiple outer NS5-branes on the gauge theory [24] with the antisymm. and conj. symm. as well as fundamentals is an open problem in the context of the analysis of supersymmetric ground states and nonsupersymmetric ground states also.

• Are there any variants of sections 2 and 3 by k NS5-branes and a single NS5'-brane? When $k = 2$, the gauge theory analysis for unitary group was given in [44]. It would be interesting to study whether there exist nonsupersymmetric meta-stable vacua and if not, how one can think of the possible deformations in the superpotential to make them stabilize at one-loop?

• When there are multiple k' middle NS5-branes, which is the same number as the one of NS5'-branes, in section 4, how one can analyze the meta-stable vacuum? For the supersymmetric brane configuration corresponding to the gauge theory [45, 46, 47], some of the analysis was done in [25] where there are the higher order term for an adjoint field in the superpotential and the interaction terms between an adjoint and quarks and symmetric tensor.

• Similarly, when there are multiple k' middle NS5-branes in section 5(there are $(2k'+1)$ middle NS5-branes in section 6), how one can analyze the meta-stable vacuum? For the supersymmetric brane configuration [29] corresponding to this gauge theory [46] which has extra two adjoint fields, the superpotential has more general form.

• It is possible to consider the case where there are multiple k NS5-branes and a single NS5'-brane(and their mirrors) in section 7. It would be interesting to study whether there exist nonsupersymmetric meta-stable vacua.

• As mentioned before, when there are multiple k' middle NS5-branes in section 8, the analysis for the supersymmetric brane configuration [25] explains the gauge theory result [46]. It is an open problem to find whether there exist nonsupersymmetric meta-stable vacua.

• Also one can think of the case when there are multiple k NS5-branes in section 9. It would be interesting to find out the supersymmetric ground states and nonsupersymmetric ground states.

• When there are multiple k' middle NS5-branes in section 10, how one can analyze the meta-stable vacuum? It is known that the supersymmetric brane configuration [25] describes the gauge theory [46]. On the other hand, one can consider the case where there exist multiple k NS5-branes with a single NS5'-brane or combination of both multiple k' middle NS5-branes and multiple k NS5-branes in section 10. For the dual of second gauge group $SU(N_c')$, we treated only $k'=1$ NS5'-brane but it is also possible to consider the general k' case where there are many meson fields.

• Also when there are multiple k NS5-branes with a single NS5'-brane, or when there are multiple k NS5-branes and multiple k' NS5'-branes in section 11, how the meta-stable vacua as well as supersymmetric ones appear?

Acknowledgments

This work was supported by grant No. R01-2006-000-10965-0 from the Basic Research Program of the Korea Science & Engineering Foundation. I would like to thank KIAS(Korea Institute for Advanced Study) for hospitality where this work was undertaken.

References

[1] K. Intriligator, N. Seiberg and D. Shih, JHEP **0604**, 021 (2006).

[2] K. Intriligator and N. Seiberg, Class. Quant. Grav. **24**, S741 (2007).

[3] H. Ooguri and Y. Ookouchi, Phys. Lett. B **641**, 323 (2006).

[4] S. Franco, I. Garcia-Etxebarria and A. M. Uranga, JHEP **0701**, 085 (2007).

[5] I. Bena, E. Gorbatov, S. Hellerman, N. Seiberg and D. Shih, JHEP **0611**, 088 (2006).

[6] A. Giveon and D. Kutasov, JHEP **0802**, 038 (2008).

[7] A. Giveon and D. Kutasov, Nucl. Phys. B **796**, 25 (2008).

[8] A. Giveon and D. Kutasov, Nucl. Phys. B **778**, 129 (2007).

[9] A. Giveon and D. Kutasov, Rev. Mod. Phys. **71**, 983 (1999).

[10] S. Elitzur, A. Giveon, D. Kutasov, E. Rabinovici and A. Schwimmer, Nucl. Phys. B **505**, 202 (1997).

[11] C. Ahn, Class. Quant. Grav. **24**, 1359 (2007).

[12] C. Ahn, Phys. Lett. B **665**, 418 (2008).

[13] C. Ahn, Phys. Lett. B **663**, 125 (2008).

[14] C. Ahn, arXiv:0801.0619 [hep-th].

[15] C. Ahn, arXiv:0802.0044 [hep-th].

[16] J. de Boer and Y. Oz, Nucl. Phys. B **511**, 155 (1998).

[17] L. Mazzucato, Y. Oz and S. Yankielowicz, JHEP **0711**, 094 (2007).

[18] A. Hanany and E. Witten, Nucl. Phys. B **492**, 152 (1997).

[19] C. Ahn, Phys. Lett. B **426**, 306 (1998).

[20] C. Ahn, K. Oh and R. Tatar, J. Geom. Phys. **28**, 163 (1998).

[21] C. Ahn, Phys. Lett. B **647**, 493 (2007).

[22] K. Landsteiner, E. Lopez and D. A. Lowe, JHEP **9807**, 011 (1998).

[23] C. Ahn, JHEP **0705**, 053 (2007).

[24] K. A. Intriligator, R. G. Leigh and M. J. Strassler, Nucl. Phys. B **456**, 567 (1995).

[25] I. Brunner, A. Hanany, A. Karch and D. Lust, Nucl. Phys. B **528**, 197 (1998).

[26] K. Landsteiner, E. Lopez and D. A. Lowe, JHEP **9802**, 007 (1998).

[27] S. Elitzur, A. Giveon, D. Kutasov and D. Tsabar, Nucl. Phys. B **524**, 251 (1998).

[28] C. Ahn, Class. Quant. Grav. **24**, 3603 (2007).

[29] J. H. Brodie and A. Hanany, Nucl. Phys. B **506**, 157 (1997).

[30] C. Ahn, Class. Quant. Grav. **25**, 075001 (2008).

[31] C. Ahn and R. Tatar, Phys. Lett. B **413**, 293 (1997).

[32] C. Ahn, arXiv:0708.4255 [hep-th].

[33] C. Ahn, arXiv:0710.0180 [hep-th].

[34] A. Giveon, A. Katz and Z. Komargodski, JHEP **0806**, 003 (2008).

[35] C. Ahn, Mod. Phys. Lett. A **22**, 2329 (2007).

[36] R. Tatar, Phys. Lett. B **419**, 99 (1998).

[37] C. Ahn, K. Oh and R. Tatar, J. Geom. Phys. **31**, 301 (1999).

[38] E. Lopez and B. Ormsby, JHEP **9811**, 020 (1998).

[39] C. Ahn, JHEP **0708**, 021 (2007).

[40] C. Ahn, Nucl. Phys. B **790**, 281 (2008).

[41] C. Ahn, Class. Quant. Grav. **25**, 095018 (2008)

[42] C. Ahn, arXiv:0711.0082 [hep-th].

[43] C. Ahn, arXiv:0705.0056 [hep-th].

[44] A. Amariti, L. Girardello and A. Mariotti, JHEP **0612**, 058 (2006).

[45] J. H. Brodie, Nucl. Phys. B **478**, 123 (1996).

[46] J. H. Brodie and M. J. Strassler, Nucl. Phys. B **524**, 224 (1998).

[47] L. Mazzucato, JHEP **0512**, 026 (2005).